



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | | |
|--|--|---|--|
| (51) International Patent Classification ⁶ : C12N 15/12, C07K 14/47 | | A3 | (11) International Publication Number: WO 99/09054 |
| | | | (43) International Publication Date: 25 February 1999 (25.02.99) |
| (21) International Application Number: PCT/BE98/00124 | | (81) Designated States: CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). | |
| (22) International Filing Date: 20 August 1998 (20.08.98) | | | |
| (30) Priority Data: 9700692 20 August 1997 (20.08.97) BE | | Published With international search report. | |
| (71) Applicants (for all designated States except US): UNIVERSITE CATHOLIQUE DE LOUVAIN [BE/BE]; Halles Universitaires, Place de l'Université 1, B-1348 Louvain-la-Neuve (BE). UNIVERSITE DE MONS-HAINAUT [BE/BE]; Place du Parc 20, B-7000 Mons (BE). | | (88) Date of publication of the international search report: 15 April 1999 (15.04.99) | |
| (72) Inventors; and (75) Inventors/Applicants (for US only): KNOOPS, Bernard [BE/BE]; Rue Chapelle Notre-Dane 3/1, B-1341 Ceroux-Mousty (BE). HERMANS, Cedric [BE/BE]; Avenue des Glycines 42, B-1030 Brussels (BE). BERNARD, Alfred [BE/BE]; Avenue de la Chapelle 6, B-1200 Brussels (BE). WATTIEZ, Ruddy [BE/BE]; Chemin du Sauvelon 17, B-7022 Hyon (BE). FALMAGNE, Paul [BE/BE]; Rue du Point du Jour 8, B-7022 Mesvin (BE). | | | |
| (74) Agents: VAN MALDEREN, Eric et al.; Office Van Malderen, Place Reine Fabiola 6/1, B-1083 Brussels (BE). | | | |

(54) Title: PEROXISOME-ASSOCIATED POLYPEPTIDE, NUCLEOTIDE SEQUENCE ENCODING SAID POLYPEPTIDE AND THEIR USES IN THE DIAGNOSIS AND/OR THE TREATMENT OF LUNG INJURIES AND DISEASES, AND OF OXIDATIVE STRESS-RELATED DISORDERS

(57) Abstract

The present invention is related to an isolated and purified polypeptide which amino acid sequence presents more than 70 % with the sequence SEQ ID NO 1. The present invention is also related to the nucleotide sequence encoding said amino acid sequence, the inhibitor directed against said sequences and their use in the diagnosis, treatment and/or prevention of lung injuries or diseases and oxidative stress-related disorders.

CLUSTAL V alignment of human and rat B18 amino acid sequences (Identity: 90%, Homology: 97.5%):

```

B18hum  MAPIKVGDALPAVEVFEGEPGNKVNLAELFKGKKGVLFGVPGAFPGCSK = SEQIDNO1
B18rat  MAPIKVGDITPSVEVFEGEPGNKVNLAELFKDKKGVLFQVPGAFPGCSK
.....
B18hum  THLPGFVEQAEALKAGQVQVACLSDVDAFVTGEWGRAHKAEGKVRLLAD
B18rat  THLPGFVEQAGALKAGQVQVACLSDVDFVTAEWGRAHQAEGKVRLLAD
.....
B18hum  PTGAFGKETDLLDDSLVSI FGNRLKRFSMVVDGIVKALNVEPDGTGL
B18rat  PTGAFGKETDLLDDSLVSL FGNRLKRFSMVIDKGIVKALNVEPDGTGL
.....
B18hum  TCSLAPNIIISQL
B18rat  TCSLAPNIIISQL
.....

```

CLUSTAL V alignment of human and mouse B18 amino acid sequences (Identity: 91%, Homology: 96%):

```

B18hum  MAPIKVGDALPAVEVFEGEPGNKVNLAELFKGKKGVLFGVPGAFPGCSK
B18mouse MAPIKVGDALPSVEVFEGEPGNKVNLAELFKGKKGVLFGVPGAFPGCSK
.....
B18hum  THLPGFVEQAEALKAGQVQVACLSDVDAFVTGEWGRAHKAEGKVRLLAD
B18mouse THLPGFVEQAGALKAGQVQVACLSDVDFVTEWGRAHQAEGKVRLLAD
.....
B18hum  PTGAFGKETDLLDDSLVSI FGNRLKRFSMVVDGIVKALNVEPDGTGL
B18mouse PTGAFGKATDLLDDSLVSL FGNRLKRFSMVIDGIVKALNVEPDGTGL
.....
B18hum  TCSLAPNIIISQL
B18mouse TCSLAPNIIISQL
.....

```

CLUSTAL V alignment of human and rat cDNA sequences (Identity: 612/780, 78.5%):

```

B18hum  GCCAGGAGGCGGAGTGGAAGTGGCCGTGGGGCGGGTATGGGACTAGCTGG
B18rat  TG-----CGTC-----CTAGGCAG
.....
B18hum  CGTGTGGCCCTGAGACGCTCAGCGGGCTATATACTCGTGGTGGGGCCG
B18rat  CATA---GCC---GGA---TCGGTGTCCGTGCATCGGCTACTTGGAC---
.....
B18hum  GCGGTCACTGTGCGGCAGCGGCAAGACGCTGCACTGAAGCAGAGTGG
B18rat  GTGCGTGGCAGGAGGAGGCGG---AAGGAGCAGGTGG
.....

```

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | | | TR | Turkey |
| BG | Bulgaria | HU | Hungary | ML | Mali | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MN | Mongolia | UA | Ukraine |
| BR | Brazil | IL | Israel | MR | Mauritania | UG | Uganda |
| BY | Belarus | IS | Iceland | MW | Malawi | US | United States of America |
| CA | Canada | IT | Italy | MX | Mexico | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NE | Niger | VN | Viet Nam |
| CG | Congo | KE | Kenya | NL | Netherlands | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NO | Norway | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | NZ | New Zealand | | |
| CM | Cameroon | | | PL | Poland | | |
| CN | China | KR | Republic of Korea | PT | Portugal | | |
| CU | Cuba | KZ | Kazakstan | RO | Romania | | |
| CZ | Czech Republic | LC | Saint Lucia | RU | Russian Federation | | |
| DE | Germany | LI | Liechtenstein | SD | Sudan | | |
| DK | Denmark | LK | Sri Lanka | SE | Sweden | | |
| EE | Estonia | LR | Liberia | SG | Singapore | | |

INTERNATIONAL SEARCH REPORT

Inter. .nal Application No

PCT/BE 98/00124

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12N15/12 C07K14/47

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|--|-----------------------|
| X | DATABASE EMBL Accssion number N42215, 27 January 1996 HILLIER L. ET AL.: "N42215 164154 hypothetical protein HI0572 - Haemophilus influenzae" XP002089366 see abstract | 8,9 |
| P,X | --- DATABASE EMBL Accession number AA639364, 1 November 1997 STRAUSBERG R.: "nq87d08.s1 NCI CGAP Co9 Homo sapiens cDNA clone IMAGE:1159311 similar to TR:G558080 G558080 PUTATIVE PEROXISOMAL PROTEIN." XP002089367 see abstract --- -/- | 8,9 |

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

7 January 1999

Date of mailing of the international search report

12.02.99

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Chakravarty, A

INTERNATIONAL SEARCH REPORT

Inter. Application No

PCT/BE 98/00124

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category ° | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|---|-----------------------|
| A | <p>DATABASE EMBL Accession number P73728, 1 February 1997 TABATA S.: "MEMBRANE PROTEIN from SYNECHOCYSTIS SP." XP002089368 see abstract</p> <p style="text-align: center;">-----</p> | 1-4 |

INTERNATIONAL SEARCH REPORT

International application No.
PCT/BE 98/00124

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 10, 11-15 (part)
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Claims Nos.: 10, 11-15 (part)

Claim 10 is directed to inhibitors of the amino acid / nucleotide sequences of claims 1-8. However, no such compounds are defined in the application. No search can be carried out for such purely speculative claims whose wording is, in fact, a mere recitation of the required result.


The same objection applies, mutatis mutandis, to claims 11-15 in as much as these claims refer to the subject-matter of claim 10.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | | |
|---|--|---|---|
| (51) International Patent Classification ⁶ : C07K 14/00 | | A2 | (11) International Publication Number: WO 99/09054 |
| | | (43) International Publication Date: 25 February 1999 (25.02.99) | |
| (21) International Application Number: PCT/BE98/00124 | | (81) Designated States: CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). | |
| (22) International Filing Date: 20 August 1998 (20.08.98) | | | |
| (30) Priority Data: 9700692 20 August 1997 (20.08.97) BE | | Published Without international search report and to be republished upon receipt of that report. | |
| (71) Applicants (for all designated States except US): UNIVERSITE CATHOLIQUE DE LOUVAIN [BE/BE]; Halles Universitaires, Place de l'Université 1, B-1348 Louvain-la-Neuve (BE). UNIVERSITE DE MONS-HAINAUT [BE/BE]; Place du Parc 20, B-7000 Mons (BE). | | | |
| (72) Inventors; and | | | |
| (75) Inventors/Applicants (for US only): KNOOPS, Bernard [BE/BE]; Rue Chapelle Notre-Dame 3/1, B-1341 Ceroux-Mousty (BE). HERMANS, Cedric [BE/BE]; Avenue des Glycines 42, B-1030 Brussels (BE). BERNARD, Alfred [BE/BE]; Avenue de la Chapelle 6, B-1200 Brussels (BE). WATTIEZ, Ruddy [BE/BE]; Chemin du Sauvelon 17, B-7022 Hyon (BE). FALMAGNE, Paul [BE/BE]; Rue du Point du Jour 8, B-7022 Mesvin (BE). | | | |
| (74) Agents: VAN MALDEREN, Eric et al.; Office Van Malderen, Place Reine Fabiola 6/1, B-1083 Brussels (BE). | | | |

(54) Title: PEROXISOME-ASSOCIATED POLYPEPTIDE, NUCLEOTIDE SEQUENCE ENCODING SAID POLYPEPTIDE AND THEIR USES IN THE DIAGNOSIS AND/OR THE TREATMENT OF LUNG INJURIES AND DISEASES, AND OF OXIDATIVE STRESS-RELATED DISORDERS



(57) Abstract

The present invention is related to an isolated and purified polypeptide which amino acid sequence presents more than 70 % with the sequence SEQ ID NO 1. The present invention is also related to the nucleotide sequence encoding said amino acid sequence, the inhibitor directed against said sequences and their use in the diagnosis, treatment and/or prevention of lung injuries or diseases and oxidative stress-related disorders.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Larvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | | | TR | Turkey |
| BG | Bulgaria | HU | Hungary | ML | Mali | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MN | Mongolia | UA | Ukraine |
| BR | Brazil | IL | Israel | MR | Mauritania | UG | Uganda |
| BY | Belarus | IS | Iceland | MW | Malawi | US | United States of America |
| CA | Canada | IT | Italy | MX | Mexico | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NE | Niger | VN | Viet Nam |
| CG | Congo | KE | Kenya | NL | Netherlands | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NO | Norway | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | NZ | New Zealand | | |
| CM | Cameroon | | | PL | Poland | | |
| CN | China | KR | Republic of Korea | PT | Portugal | | |
| CU | Cuba | KZ | Kazakstan | RO | Romania | | |
| CZ | Czech Republic | LC | Saint Lucia | RU | Russian Federation | | |
| DE | Germany | LI | Liechtenstein | SD | Sudan | | |
| DK | Denmark | LK | Sri Lanka | SE | Sweden | | |
| EE | Estonia | LR | Liberia | SG | Singapore | | |

1430 Rec'd PCT/PTO 22 FEB 2000

5

10 PEROXISOME-ASSOCIATED POLYPEPTIDE, NUCLEOTIDE SEQUENCE
ENCODING SAID POLYPEPTIDE AND THEIR USES IN THE DIAGNOSIS
AND/OR THE TREATMENT OF LUNG INJURIES AND DISEASES, AND OF
OXIDATIVE STRESS-RELATED DISORDERS

Field of the invention

15 The present invention is related to a new
peroxisome-associated polypeptide, the nucleotide sequence
encoding said polypeptide and portions thereof as well as
their uses in the diagnosis of several diseases, especially
the diagnosis and/or the treatment of lung injuries and
20 diseases, and of oxidative stress-related disorders.

Background of the invention

The peroxisomes are organelles nearly
ubiquitous in eukaryotic cells. They contain enzymes
25 essential for various catabolic and anabolic pathways. Some
of these enzymes are expressed constitutively while others
can be induced under appropriate conditions. Peroxisomes
carry out a variety of essential reactions such as
peroxisomal oxidation and respiration, fatty acid beta-
30 oxidation, cholesterol and dolichol metabolism, ether-
phospholipid synthesis, and glyoxylate and pipercolic acid
metabolism.

The peroxisomal respiratory pathway is based upon the formation of hydrogen peroxide by a collection of oxidases and the decomposition of the H_2O_2 by catalase. These reactions are responsible for 20% of oxygen
5 consumption in liver, and several oxidases have been identified in peroxisomes. Ethanol elimination via catalase in peroxisomes may be significant in addition to the oxidation via cytosolic alcohol dehydrogenase.

The peroxisomal beta-oxidation system
10 catalyses the beta-oxidative chain shortening of a specific set of compounds which can not be handled by mitochondria : very long chain fatty acids, di- and trihydroxycholestanoic acids, pristanic acid, long chain dicarboxylic acids, several prostaglandins, several leukotrienes, 12- and 15-
15 hydroxyeicosatetraenoic acid, and several mono- and polyunsaturated fatty acids, which are of direct diagnostic relevance for some peroxisomal disorders.

Peroxisomes play also a major role in the synthesis of cholesterol and other isoprenoids. Fibroblasts
20 from patients affected by disorders of peroxisome biogenesis show low capacity to synthesise cholesterol.

Two enzyme activities responsible for introduction of the characteristic ether linkage in ether-linked phospholipids (dihydroacetonephosphate
25 acyltransferase (DHAPAT) and alkyl-dihydroxyacetonephosphate synthase (alkyl-DHAP synthase)) are localised in peroxisomes. These enzymes are not yet cloned. As demonstrated by the identification of patients with deficiency of either DHAPAT or alkyl-DHAP synthase with
30 severe clinical abnormalities, ether-phospholipids are of major importance in humans.

Peroxisomes are able to detoxify glyoxylate via alanine/glyoxylate aminotransferase. The deficiency of this cloned enzyme causes hyperoxaluria type I.

L-pipecolate is a minor metabolite of L-lysine and is
5 catabolised by the L-pipecolate oxidase localised in peroxisomes. The enzyme is deficient in cerebro-hepato-renal (Zellweger) syndrome.

In human, the importance of peroxisomes was emphasised by a number of inherited diseases involving
10 either a defect in the biogenesis of peroxisomes or a deficiency of one (or more) peroxisomal enzymes. So far, 12 different peroxisomal disorders have been described and most of them are lethal.

A wide variety of chemicals have been showed
15 to produce peroxisome proliferation and induction of peroxisomal and microsomal fatty acids-oxidising enzymes activities in rats and mice. Several peroxisomes proliferators have been shown to increase the incidence of liver tumours in these species. Proposed mechanisms of
20 liver tumour formation by peroxisomes proliferators include induction of sustained oxidative stress.

Therefore, newly identified molecules associated with peroxisomes could be used for the development of diagnostic tools and possibly for the
25 improvement of several therapeutical applications of various diseases associated with peroxisomal disorders. In addition, it is useful to identify the molecules present in specific organs like the lung and which may be used as specific markers of inflammatory diseases as well as lung
30 injuries or diseases.

Summary of the invention

The Inventors have isolated and purified a new sequence of a low molecular weight human broncho-alveolar polypeptide. Said mammal, preferably human, protein or polypeptide (hereafter identified as B18hum
5 protein) has been sequenced and its corresponding genomic DNA (SEQ ID NO 8) and cDNA (SEQ ID NO 1) have been identified. Similarly, the corresponding nucleotide and amino acid sequence from a rat (SEQ ID NO 3 and 4) and from
10 a mouse (SEQ ID NO 5 and 6) have been obtained.

Said sequences present several homologies with other peroxisomal proteins of yeast and comprise a carboxy-terminal tripeptide SQL which is necessary for the specific targeting and translocation of several proteins
15 into the peroxisome.

Therefore, the present invention is related to a new isolated and purified polypeptide sequence having a amino acid sequence which presents more than 70% homology, advantageously more than 85% homology, more
20 preferably more than 95% homology, with the amino acid sequence SEQ ID NO 2.; Said amino acid sequence is advantageously obtained from a mammal, preferably from a rat, a mouse or a human.

The present invention is also related to the isolated and purified polypeptide sequence corresponding to
25 the amino acid sequence SEQ ID NO 2 or a portion thereof, preferably an immunoreactive portion (putative immunogenic domain or T or B cell epitopes).

Said portions are advantageously comprised
30 between :

- Glutamic acid position 13 - Glutamic acid position 27
- Alanine position 26 - Leucine position 36

- Alanine position 42 - Glutamic acid position 57
- Glutamic acid position 57 - Valine position 69
- Valine position 80 - Leucine position 97
- Arginine position 95 - Leucine position 112
- 5 - Serine position 118 - Serine position 129
- Valine position 137 - Threonine position 150

Preferably, said portion has more than 10, 20, 30, 50 or 70 amino acids. Specific portions of the amino acid sequence SEQ ID NO 2 are also portions of more
10 than 70 amino acids which present at least 80% of the proteinic activity (see example 5) of the complete SEQ ID NO 2 sequence. Therefore, the amino acid sequence according to the invention can be partially deleted while maintaining its activity, preferably its anti-oxidative activity, which
15 will be described hereafter.

According to the invention, the amino acid sequence SEQ ID NO 2 presents a pI of 7.16 and a molecular weight of 17047 Dalton as hereafter defined by bidimensional electrophoresis.

20 The present invention is also related to the nucleotide sequence encoding the amino acid sequence according to the invention and its regulatory sequences upstream said coding sequence. A nucleotide sequence encoding the polypeptide according to the invention is a
25 genomic DNA (see SEQ ID NO 10), a cDNA (see SEQ ID NO 1) or a mRNA, possibly comprising said upstream regulatory sequence. Advantageously, said nucleotide sequence presents more than 70%, advantageously more than 85%, more preferably more than 95% homology with SEQ ID NO 1 or its
30 complementary strand.

According to a preferred embodiment of the present invention, said nucleotide sequence corresponds to the nucleotide sequence SEQ ID NO 1, its complementary strand or a portion thereof.

5 "A portion of the nucleotide sequence SEQ ID NO 1" means any nucleotide sequence of more than 15 base pairs (such as a primer, a probe or an antisense nucleotide sequence) which allow the specific identification, reconstitution or blocking of the complete nucleotide
10 sequence SEQ ID NO 1 (including its regulatory sequences upstream the coding sequence).

Said portions allow the specific identification, reconstitution or blocking by specific hybridisation with the nucleotidic sequence SEQ ID NO 1,
15 preferably under standard stringent conditions, with sequences like probes or primers possibly labelled with a compound (radioactive compound, enzyme, fluorescent marker, etc.), and can be used in a specific diagnostic or dosage method like probe hybridisation (see Sambrook et al., §§
20 9.47-9.51 in *Molecular Cloning : A Laboratory Manual*, Cold Spring Harbor, Laboratory Press, Cold Spring Harbor, New York (1989)), genetic amplification (like PCR (US patent 4,683,195), LCR (Wu et al., *Genomics* 4, pp. 560-569), CPR (US patent 5,011,769)).

25 Exemplary stringent hybridisation conditions are as follows : hybridisation at 42 °C in 50% formamide 5x SSC, 20 mM sodium phosphate, pH 6.8 washing in 0.2x SSC at 55 °C. It is understood by those skilled in the art that variation of these conditions occur based on the length and
30 GC nucleotide content of the sequence to be hybridised. Formulas standard in the art are appropriated for

determining exact hybridisation conditions (see Sambrook et al.

Preferred examples of said nucleotide portions are as follows :

| | <u>Sequence</u> | <u>Position</u> |
|----|---------------------------------|------------------------|
| 5 | 5'-gccatcccagcagtgagggtgttg-3' | (SEQ ID NO 11) 217-241 |
| | 5'-ttgaacagctctgccagggttcacc-3' | (SEQ ID NO 12) 261-284 |
| | 5'-tggaggtgtttgaaggggagccag-3' | (SEQ ID NO 13) 230-253 |
| | 5'-cagggttcaccttgttccctggctc-3' | (SEQ ID NO 14) 247-270 |
| 10 | 5'-gggtatgggactagctggcg-3' | (SEQ ID NO 15) 33-52 |
| | 5'-ctggccaacattccaattgcag-3' | (SEQ ID NO 16) 747-768 |

and the sequences of respectively 601 (SEQ ID NO 8), 604 (SEQ ID NO 9) and 469 (SEQ ID NO 7) base pairs corresponding to specific mRNA alternative splicing of the B18 human nucleotide sequence as described in Figure 4 (the known genomic sequence incorporating several introns and exons is represented in the sequence SEQ ID NO 10).

Said sequences may be used for a genetic amplification or a probe hybridisation as above-described.

The present invention is also related to a vector comprising the necessary elements for the injection, transfection or transduction of cells and having incorporated one or more of the nucleotide sequences according to the invention. The vector according to the invention is selected from the group consisting of viruses, plasmids, phagemides, cationic vesicles, liposomes or a mixture thereof. Said vector may comprise also one or more adjacent regulatory sequences (such as promoter(s), secretion and termination signal sequence(s)), advantageously operably linked to the nucleotide sequence according to the invention.

The present invention is also related to the cell transformed by said vector and expressing the polypeptide according to the invention.

The nucleotide sequence according to the invention can be also introduced in said cell by the formation of CaPO_4 -nucleic acid precipitate, DEAE-dextran-nucleic acid complex or by electroporation.

Another aspect of the present invention is related to an inhibitor of the polypeptide according to the invention or the nucleotide sequence according to the invention (including the upstream sequences like promoter-operator regulatory sequence which may be inhibited by a cis- and/or transactivating repressor). Said inhibitor is advantageously an antibody or a fragment of said antibody such as an hypervariable portion of said antibody directed against the amino acid or nucleotide sequence of the polypeptide according to the invention. Other examples of inhibitors according to the invention are antisense nucleotide sequences which allow the blocking of the expression of the nucleotide sequence according to the invention.

Another aspect of the present invention is related to a diagnostic device (such as a diagnostic kit or a chromatographic column) comprising an element selected from the group consisting of the amino acid sequence of said polypeptide, its nucleotide sequence, and/or the inhibitor according to the invention or a fragment thereof as above-described. Said diagnostic device may comprise also necessary reactants and media for the diagnostic and/or dosage of the nucleotide and/or amino acid sequence of the polypeptide according to the invention, which are based upon the method selected from the group consisting of

in situ hybridisation, hybridisation by labelled antibodies, especially RIA (Radio Immuno Assay) or ELISA (Enzymes Linked Immuno-Sorbent Assay) technologies, detection upon filter, upon solid support, in solution, in sandwich, upon gel, dot blot hybridisation, Northern blot hybridisation, Southern blot hybridisation, isotopic or non-isotopic labelling (by immunofluorescence or biotinilised probes), genetic amplification, (especially by PCR or LCR), double immunodiffusion technique, counter-electrophoresis technique, haemagglutination or a mixture thereof.

Another aspect of the present invention concerns a diagnosis method wherein a biological sample from the patient, such as cephalo-rachidian fluid, serum, blood, plasma, urine, broncho-alveolar lavage, stomach lavage, etc., is isolated from the patient, and is put in contact with the diagnostic device according to the invention for the diagnosis or the monitoring of an injury or a disease, preferably a lung injury or an oxidative stress-related disorder, affected by the presence of pro-oxidant agent or oxidative stress such as specific cardiovascular diseases like arteriosclerosis, neurodegenerative disorders (Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis), apoptosis, inflammatory reactions, allergic reactions such as asthma, hay fever and eczema, high bone mass syndrome, osteopetrosis, osteoporosis-pseudoglioma syndrome, and Bardet-Biedl syndrome 1. Said diagnosis and monitoring upon one or more biological samples obtained from several tissues from the patient can be advantageously obtained by one or more of the methods above-described, which could be adapted

according to the specific biological sample by the person skilled in the art.

Therefore, the product according to the invention could be used as a marker for the above-
5 identified injuries, diseases or disorders in a broad spectrum of tissues as shown in the enclosed Figure 1.

A further aspect of the present invention is related to a pharmaceutical composition comprising a pharmaceutically acceptable carrier and an element selected
10 from the group consisting of the nucleotide sequence, the amino acid sequence of the polypeptide according to the invention, the inhibitor directed against said sequences and/or one or more portions thereof.

A last aspect of the present invention is
15 related to the use of the pharmaceutical composition according to the invention for the manufacture of a medicament in the treatment and/or the prevention of lung injuries and/or diseases or of oxidative stress-related disorders.

20 The present invention is also related to a prevention and/or treatment method of a patient, especially a human patient, preferably affected by lung injuries and/or diseases or by oxidative stress-related disorders, wherein a sufficient amount of the pharmaceutical
25 composition according to the invention is administered to said patient in order to treat, avoid and/or reduce the symptoms of said injuries and/or diseases.

Other injuries and/or diseases which can be prevented and/or treated are injuries and/or diseases
30 affected by the presence of pro-oxidant agents or oxidative stress, such as specific cardio-vascular diseases like arteriosclerosis, neurodegenerative disorders such as

Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, apoptosis and inflammatory reactions and some allergic reactions such as asthma, hay fever and eczema, high bone mass syndrome, osteopetrosis, 5 osteoporosis-pseudoglioma syndrome, and Bardet-Biedl syndrome 1.

The pharmaceutically acceptable carrier according to the invention is any compatible non-toxic substance suitable for administering the composition 10 according to the invention to a human patient. Pharmaceutically acceptable carriers according to the invention suitable for oral administration are the ones well known by the person skilled in the art, such as tablets, coated or non-coated pills, capsules, spray-gas, 15 patches, gels, solutions or syrups. Pharmaceutically acceptable carriers vary according to the mode of administration (intravenous, intramuscular, subcutaneous, parenteral, etc.), and may comprise also adjuvants well known by the person skilled in the art to increase, reduce 20 and/or regulate humoral, local and/or cellular response of the immune system.

The pharmaceutical composition according to the invention may be prepared by the methods, generally applied by the person skilled in the art in the preparation 25 of various pharmaceutical compositions, wherein the percentage of the active compound/pharmaceutically acceptable carrier can vary within very large ranges, only limited by the tolerance of the patient to said pharmaceutical composition, and wherein the limits are 30 particularly determined by the frequency of administration and the possible side-effects of the active compounds or its pharmaceutically acceptable carrier.

Another aspect of the invention is related to the use of the diagnostic device according to the invention for performing upon the patient or upon a biological fluid obtained from the patient, a diagnosis, a dosage and/or a
5 monitoring of the above-mentioned injuries or diseases or oxidative stress-related disorders affecting the patient.

A further aspect of the present invention is related to a cell or a non-human animal, preferably a mammal such as a mouse or a rat, transformed by the vector
10 according to the invention and overexpressing the polypeptide according to the invention, or a non-human animal, preferably a mammal such as a mouse or a rat, genetically modified by a partial or total deletion of its genomic sequence encoding the polypeptide according to the
15 invention (knock-out non-human mammal) and obtained by methods well known by the person skilled in the art such as the one described by Kahn et al. (*Cell*, Vol. 92, pp. 593-596 (March 1998)).

Other examples of genetically modified non-
20 human animals according to the invention may be a transgenic non-human animal comprising an inhibitor according to the invention, preferably an antisense nucleic acid sequence complementary to the nucleotide sequence according to the invention so placed as to be transcribed
25 into antisense mRNA which is complementary to the nucleotide sequence according to the invention and which hybridises to said nucleotide sequence, thereby reducing or blocking its translation.

Further aspects of the present invention will
30 be described in the enclosed non-limiting examples in reference to the following Figures.

Brief description of the drawings

- Figure 1 represents a dot blot analysis of mRNA encoding the polypeptide according to the invention in various types of human tissues.
- 5 Figure 2 represents a Northern blot analysis of mRNA encoding the polypeptide according to the invention in a rat lung after administration of lipopolysaccharides (LPS) inducing an inflammatory reaction of the lung.
- 10 Figure 3 represents a Northern blot analysis of mRNA encoding the polypeptide according to the invention in a rat lung after intraperitoneal injection of pneumotoxicants.
- Figure 4 is a schematic representation of the human genomic sequence, the complete cDNA sequence and the corresponding amino acid sequence.
- 15 Figure 5 represents respectively the alignment of the sequences of the human B18 polypeptide according to the invention with the corresponding rat and mouse sequences.
- 20

Example 1 : Homology between the B18 polypeptide according to the invention with other known nucleotide or amino acid sequences

- 25 The BLAST 2.0 software (gapped BLAST at the NCBI Internet site) was used for searching for homologies between human B18 (162 amino acids) and known polypeptides in databases (GenBank, SwissProt). Said search did not give perfect alignment with known peptides from different
- 30 species (Table 1). Homologies of the human B18 cDNA (805 nucleotides) with GenBank, EMBL, DDBJ and PDB deposited

nucleotide sequences (Table 2) and GenBank Expression Sequence TAGS (ESTs) were noted.

Table 1 : Homologies of the B18 proteins (162 amino acid) with other proteins

| Name | NCBI ID | Identity (%) Homology (%) |
|--|----------|------------------------------|
| Membrane protein (synechocystis sp.) | 1652859 | 57/129 (44%) 81/129 (62%) |
| Peroxisomal-like protein (Aspergillus fumigatus) | 2769700 | 56/176 (31%) 90/176 (50%) |
| Haein HI0572 hypothetical protein (Haemophilus influenzae) | 1723174 | 53/146 (36%) 80/146 (54%) |
| PMP20 (Schizosaccharomyces pombe) | AJ002536 | 54/161 (33%) 85/161 (52%) |
| Peroxisomal membrane protein A (PMP 20) (Candida boidinii) | 130360 | 59/170 (34%) 89/170 (51%) |
| Peroxisomal membrane protein B (PMP 20) (Candida boidinii) | 130361 | 58/170 (34%) 88/170 (51%) |
| Putative peroxisomal protein PMP from yeast (Saccharomyces cerevisiae) | 1709682 | 41/138 (29%) 72/138 (51%) |
| Alkylhydroperoxide reductase C22 protein (Escherichia coli) | P26427 | 36/126 (28%) 58/126 (45%) |

Table 2

| Name | Access NO | Identity |
|--|-----------|---------------|
| Human mRNA down-regulated in cells infected by adenovirus 5 | U82616 | 259/263 (98%) |
| Human mRNA down-regulated in cells infected by adenovirus 5 | U82615 | 300/321 (93%) |

In the Table 2, an identity of 98% has been obtained with the alignment of 259 nucleotides of CDNA B18, which comprises in its totality 805 nucleotides, with 263 nucleotides of U82616 CDNA. A similar identity has been
5 obtained with the U82615 sequence.

The sequence SEQ ID NO 1 comprising 805 nucleotides presents a homology with several EST sequences obtained from a human and from a mouse, having the following references :

10 Human :

AA130751, N42215, W38597, N91311, N68467, AA187737, N68916, W00593, R88950, AA181884, H20154, H66666

Mouse :

AA220019, AA123351, AA087129, AA255021, AA249897, W71344
15

Example 2 : Tissue detection

A human RNA master Blot (Clontech) containing 100-500 ng of poly-A + human RNA in each dot (normalised to the mRNA expression levels of eight different housekeeping
20 genes) was hybridised with a 554 bp-long B18 probe labelled with ^{32}P , and quantified, using Phosphorimaging Technology. As shown in Figure 1, B18 mRNA is present in all tissues examined but predominantly in trachea, lung, kidney, thyroid gland, stomach, colon, heart and some regions of
25 the brain. Highest expression has been noted in the thyroid tissue. This presence is probably correlated with the possible antioxidant activity of the B18 polypeptide according to the invention.

30 Example 3 : Inflammatory reaction

Figure 2 represents a Northern blot analysis of rat lung mRNA after 6, 48 and 72 hours after

lipopolysaccharides (LPS) instillation inducing an inflammatory reaction in the lung.

A Northern blot containing 15 μ g of total RNA in each lane was hybridised with a 225 bp-long rat B18 probe, stripped and reprobed with a 572 bp-long rat β -actin probe, both labelled with 32 P. Northern blot was quantified using Phosphorimaging Technology and the B18 mRNA data were normalised to β -actin mRNA level.

10 Example 4 : Pneumotoxic reaction

Figure 3 represents a Northern blot analysis of rat lung mRNA after intraperitoneal injection of pneumotoxicants (4-ipomeanol, 1-(3-furyl)-4-hydroxypentanone (IPO), methylcyclopentadienyl manganese tricarbonyl (MMT) or alpha naphthylthiourea (ANTU)). These agents are known to induce in the lung acute lesions of Clara (IPO) and alveolar cells (MMT) as well as increasing the permeability of the alveolar/blood barrier (ANTU). A Northern blot containing 15 μ g of total RNA in each lane was hybridised with a 225 bp-long rat B18 probe, stripped and reprobed with a 572 bp-long β -actin probe both labelled with 32 P. The Northern blot was quantified using Phosphorimaging Technology and rat B18 mRNA data were normalised to β -actin mRNA level.

25

Example 5 : Proteinic activity of the B18 polypeptide

An amino analysis of the complete human B18 amino acid sequence shows that said polypeptide presents specific portions showing an homology with other anti-oxidant enzymes (starting from a Leucine at position 36 until a Cysteine at position 47) and an other portion

having an important homology with beta chains of ATP synthase (starting from a Glutamic acid at position 13 until a Glycine in position 38).

Furthermore, the B18 amino acid sequence
5 according to the invention shows an important homology with an *Aspergillus fumigatus* allergen (34% identity and 60% homology by using clustal V sequence alignment), especially upon the portion of said B18 polypeptide having possible antioxidant properties. Therefore, it is possible that a
10 peroxisomal protein (possibly homologous to B18 protein) is able to induce and to bind IgE from patients sensitised to *Aspergillus fumigatus* peroxisomal proteins after an induction of the patient immune system with *Aspergillus fumigatus* allergen. This mechanism can be compared to a
15 reaction obtained with the manganese superoxide dismutase (MnSOD) wherein the human MnSOD is able to bind to IgE from patients sensitised to *Aspergillus fumigatus* MnSOD.

Furthermore, the Inventors have identified a portion of the B18 human polypeptide which presents an
20 homology with a Cyclophilin-binding domain of *Candida boidinii* PMP20 (receptor, of the immuno-suppressant drug cyclosporine A). Said possible Cyclophilin-binding domain is starting from the Threonine in position 150 until the Leucine in position 161.

25

Example 6 : B18 human gene and mRNA alternative splicing

As represented in the enclosed Figure 4, the Inventors have identified upon the genomic DNA (SEQ ID NO 10) 5 exons and 5 introns. By RT-PCR (using primers 5'-
30 gggatatgggactagctggcg-3' and 5'-ctggccaacattccaattgcag-3') and according to the genomic sequence, 4 different cDNAs corresponding to the transcription of the said genomic DNA

have been identified in human lung and in human brain. A first cDNA of 736 bp corresponds to the cDNA encoding the complete amino acid sequence of the B18 protein according to the invention. However, 3 other cDNAs of 601, 604 and 5 469 bp were also identified, and comprise specific splicings of one or more exons.

Therefore, another aspect of the present invention is related to said specific portions of the complete genomic or CDNA nucleotide sequence according to 10 the invention or to specific portions of the complete amino acid sequence of the B18 protein according to the invention, which could be used also as specific markers of the B18 activity, preferably the anti-oxidative activity.

15 Example 7 : Knock-out mouse

Exons of a mouse genomic sequence encoding the B18 polypeptide according to the invention have been deleted by homologous recombination. Said homologous recombination has been obtained with a genetic sequence 20 comprising a neomycin resistant gene. The targeting vector with said gene and a ,thymidine kinase (in order to eliminate non-homologous recombinants with ganciclovir) has been prepared. Said recombination was used for the deletion of one or more exons of the B18 polypeptide. After 25 electroporation of ES cells with the targeting vector, positive clones having incorporated homologous recombination were identified by Southern blot with labelled probes. Aggregation of said positive clones with a morula from a Swiss pseudo-pregnant mouse produces several 30 chimeric mice which survive after birth. Several homozygote mice are obtained by cross-breeding and are used as a model for the above-mentioned diseases.

Similar experiments may be done with another mammal whose B18 sequence is known (the B18 sequence of a mouse and a rat and their alignment with the human sequence is shown in the enclosed Figure 5).

5

Example 8 : Chromosome localisation of human B18 gene

Radiation hybrid clones (GeneBridge 4 Radiation Hybrid Panel, Research Genetics) were used for performing chromosome localisation by PCR with two
10 different pairs of primers (5'-caggttcaccttggtccctggctc-3' (SEQ ID NO 14), 5'-atgttatgcaaccctttgcgacac-3' (SEQ ID NO 17) and 5'-gtgtttgaaggggagccaggaac-3' (SEQ ID NO 18), 5'-agagacagggtttcaccatcttgg-3' (SEQ ID NO 19)).

The Inventors have located B18 genomic
15 sequence on human chromosome 11q13. B18 gene has been located 7.15-6.1 cR from marker D11S913 between markers D11S1963 and D11S4407 (Genome Database internet site).

Unknown genes linked to different disorders have been localised in the same region of chromosome 11.
20 Therefore, B18 gene is possibly associated with these disorders:

- atopy (atopic hypersensitivity: asthma, hay fever and eczema; MIM n°147050 at OMIM of NCBI internet site),
- high bone mass syndrome (MIM n°601884),
- 25 - osteopetrosis (MIM n°259700),
- osteoporosis-pseudoglioma syndrome (MIM n°259770) and
- Bardet-Biedl syndrome 1 (MIM n°209901).

CLAIMS

1. Amino acid sequence having more than 70% homology with the sequence SEQ ID NO 2.

2. Amino acid sequence according to claim 1,
5 having more than 85% homology with the sequence SEQ ID NO 2.

3. Amino acid sequence according to claim 1 or 2, having more than 95% homology with the sequence SEQ ID NO 2.

10 4. Amino acid sequence according to any one of the preceding claims, corresponding to SEQ ID NO 2 or an immunoreactive portion thereof.

5. Nucleotide sequence encoding the amino acid sequence according to any one of the preceding claims
15 and presenting more than 70% homology with SEQ ID NO 1 or its complementary strand.

6. Nucleotide sequence according to claim 5, having more than 85% homology with the sequence SEQ ID NO 1 or its complementary strand.

20 7. Nucleotide sequence according to claim 5 more than 95% homology with the sequence SEQ ID NO 1 or its complementary strand.

8. Nucleotide sequence according to any one of the claims 5 to 7, corresponding to the sequence SEQ ID
25 NO 1, its complementary strand or a portion thereof specific for SEQ ID NO 1 and comprising more than 15 base pairs.

9. Vector comprising the nucleotide sequence according to any one of the claims 5 to 8.

30 10. Inhibitor directed against the amino acid or nucleotide sequence according to any one of the claims 1 to 8.

11. Inhibitor according to claim 10, being an antibody, preferably a monoclonal antibody, or a portion of said antibody.

12. Diagnostic device comprising an element
5 selected from the group consisting of the amino acid sequence according to any one of the claims 1 to 4, the nucleotide sequence according to any one of the claims 5 to 8, the inhibitor according to claim 10 or 11, their portions or a mixture thereof.

10 13. Method for the in vitro detection of lung injuries and diseases or oxidative stress-related diseases and disorders, especially inflammatory diseases, comprising the steps of :

- isolating a sample from a body fluid of a patient,
15 preferably a human patient,
- possibly inhibiting the contaminants present in said sample,
- put in contact said sample with an element selected from the group consisting of the amino acid sequence
20 according to any one of the claims 1 to 4, the nucleotide sequence according to any one of the claims 5 to 8, the inhibitor according to claim 10 or 11, their portions or a mixture thereof, and
- detecting a reaction of a molecule present in said
25 sample with said element.

14. Pharmaceutical composition comprising a pharmaceutically acceptable carrier and an element selected from the group consisting of the amino acid sequence according to any one of the claims 1 to 4, the nucleotide
30 sequence according to any one of the claims 5 to 8, the

inhibitor according to claim 10 or 11, their portions or a mixture thereof.

15 15. Use of the pharmaceutical composition according to claim 14 for the manufacture of a medicament for the prevention and/or the treatment of lung injuries or diseases, and of oxidative stress-related diseases or disorders, such as specific cardio-vascular diseases like arteriosclerosis, neurodegenerative disorders such as Alzheimer's disease, Parkinson's disease, amyotrophic
10 lateral sclerosis, apoptosis and inflammatory reactions, allergic reactions such as asthma, hay fever and eczema, high bone mass syndrome, osteopetrosis, osteoporosis-pseudoglioma syndrome, and Bardet-Biedl syndrome 1.

15 16. Cell transformed by the vector according to claim 9 or comprising a partial or total deletion of its nucleotide sequence according to any one of the claims 5 to 8.

20 17. Non-human animal, preferably a mammal, transformed by the vector according to claim 9 or comprising a partial or total deletion of its nucleotide sequence according to any,one of the claims 5 to 8.

1
SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: UNIVERSITE CATHOLIQUE DE LOUVAIN
Halles Universitaires
(B) STREET: Place de l' Universite, 1
(C) CITY: LOUVAIN-LA-NEUVE
(E) COUNTRY: BELGIUM
(F) POSTAL CODE (ZIP): B-1348

(A) NAME: UNIVERSITE DE MONS-HAINAUT
(B) STREET: Place du Parc 20
(C) CITY: MONS
(E) COUNTRY: BELGIUM
(F) POSTAL CODE (ZIP): B-7000

(ii) TITLE OF INVENTION: PEROXISOME-ASSOCIATED PEPTIDE, NUCLEOTIDE
SEQUENCE ENCODING SAID PEPTIDE AND THEIR USES IN THE
DIAGNOSTIC AND/OR THE TREATMENT OF LUNG INJURIES AND
DISEASES, AND OF OXIDATIVE STRESS-RELATED DISORDERS

(iii) NUMBER OF SEQUENCES: 19

(iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 805 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 193..681

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

GCCAGGAGGC GGAGTGGGAAG TGGCCGTGGG GCGGGTATGG GACTAGCTGG CGTGTGCGCC 60
CTGAGACGCT CAGCGGGGCTA TATACTCGTC GGTGGGGCCG GCGGTCAGTC TCGGCAGCG 120



2

| | |
|--|-----|
| GCAGCAAGAC GGTGCAGTGA AGGAGAGTGG GCGTCTGGCG GGGTCCGCAG TTTCAGCAGA | 180 |
| GCCGCTGCAG CC ATG GCC CCA ATC AAG GTG GGA GAT GCC ATC CCA GCA Met Ala Pro Ile Lys Val Gly Asp Ala Ile Pro Ala. 1 5 10 | 228 |
| GTG GAG GTG TTT GAA GGG GAG CCA GGG AAC AAG GTG AAC CTG GCA GAG Val Glu Val Phe Glu Gly Glu Pro Gly Asn Lys Val Asn Leu Ala Glu 15 20 25 | 276 |
| CTG TTC AAG GGC AAG AAG GGT GTG CTG TTT GGA GTT CCT GGG GCC TTC Leu Phe Lys Gly Lys Lys Gly Val Leu Phe Gly Val Pro Gly Ala Phe 30 35 40 | 324 |
| ACC CCT GGA TGT TCC AAG ACA CAC CTG CCA GGG TTT GTG GAG CAG GCT Thr Pro Gly Cys Ser Lys Thr His Leu Pro Gly Phe Val Glu Gln Ala 45 50 55 60 | 372 |
| GAG GCT CTG AAG GCC AAG GGA GTC CAG GTG GTG GCC TGT CTG AGT GTT Glu Ala Leu Lys Ala Lys Gly Val Gln Val Val Ala Cys Leu Ser Val 65 70 75 | 420 |
| AAT GAT GCC TTT GTG ACT GGC GAG TGG GGC CGA GCC CAC AAG GCG GAA Asn Asp Ala Phe Val Thr Gly Glu Trp Gly Arg Ala His Lys Ala Glu 80 85 90 | 468 |
| GGC AAG GTT CGG CTC CTG GCT GAT CCC ACT GGG GCC TTT GGG AAG GAG Gly Lys Val Arg Leu Leu Ala Asp Pro Thr Gly Ala Phe Gly Lys Glu 95 100 105 | 516 |
| ACA GAC TTA TTA CTA GAT GAT TCG CTG GTG TCC ATC TTT GGG AAT CGA Thr Asp Leu Leu Leu Asp Asp Ser Leu Val Ser Ile Phe Gly Asn Arg 110 115 120 | 564 |
| CGT CTC AAG AGG TTC TCC ATG GTG GTA CAG GAT GGC ATA GTG AAG GCC Arg Leu Lys Arg Phe Ser Met Val Val Gln Asp Gly Ile Val Lys Ala 125 130 135 140 | 612 |
| CTG AAT GTG GAA CCA GAT GGC ACA GGC CTC ACC TGC AGC CTG GCA CCC Leu Asn Val Glu Pro Asp Gly Thr Gly Leu Thr Cys Ser Leu Ala Pro 145 150 155 | 660 |
| AAT ATC ATC TCA CAG CTC TGA GGCCCTGGGC CAGATTACTT CCTCCACCCC Asn Ile Ile Ser Gln Leu * 160 | 711 |
| TCCCTATCTC ACCTGCCCAG CCCTGTGCTG GGGCCCTGCA ATTGGAATGT TGGCCAGATT | 771 |
| TCTGCAATAA ACACTTGTGG TTTGCGGAAA AAAA | 805 |

(2) INFORMATION FOR SEQ ID NO: 2:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 163 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

3

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Met | Ala | Pro | Ile | Lys | Val | Gly | Asp | Ala | Ile | Pro | Ala | Val | Glu | Val | Phe |
| 1 | | | | 5 | | | | | 10 | | | | | 15 | |
| Glu | Gly | Glu | Pro | Gly | Asn | Lys | Val | Asn | Leu | Ala | Glu | Leu | Phe | Lys | Gly |
| | | | 20 | | | | | 25 | | | | | 30 | | |
| Lys | Lys | Gly | Val | Leu | Phe | Gly | Val | Pro | Gly | Ala | Phe | Thr | Pro | Gly | Cys |
| | | 35 | | | | | 40 | | | | | 45 | | | |
| Ser | Lys | Thr | His | Leu | Pro | Gly | Phe | Val | Glu | Gln | Ala | Glu | Ala | Leu | Lys |
| | 50 | | | | | 55 | | | | | 60 | | | | |
| Ala | Lys | Gly | Val | Gln | Val | Val | Ala | Cys | Leu | Ser | Val | Asn | Asp | Ala | Phe |
| 65 | | | | | 70 | | | | | 75 | | | | 80 | |
| Val | Thr | Gly | Glu | Trp | Gly | Arg | Ala | His | Lys | Ala | Glu | Gly | Lys | Val | Arg |
| | | | | 85 | | | | | 90 | | | | | 95 | |
| Leu | Leu | Ala | Asp | Pro | Thr | Gly | Ala | Phe | Gly | Lys | Glu | Thr | Asp | Leu | Leu |
| | | | 100 | | | | | 105 | | | | | 110 | | |
| Leu | Asp | Asp | Ser | Leu | Val | Ser | Ile | Phe | Gly | Asn | Arg | Arg | Leu | Lys | Arg |
| | | 115 | | | | | 120 | | | | | 125 | | | |
| Phe | Ser | Met | Val | Val | Gln | Asp | Gly | Ile | Val | Lys | Ala | Leu | Asn | Val | Glu |
| | 130 | | | | | 135 | | | | | 140 | | | | |
| Pro | Asp | Gly | Thr | Gly | Leu | Thr | Cys | Ser | Leu | Ala | Pro | Asn | Ile | Ile | Ser |
| 145 | | | | | 150 | | | | | 155 | | | | | 160 |
| Gln | Leu | * | | | | | | | | | | | | | |

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 780 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Rattus Rattus

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 136..624

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

| | |
|---|-----|
| TGCGTCCTAG GCAGCATAGC CGGATCGGTG CTCCGTGCAT CGGCTACTTG GACGTGCGTG | 60 |
| GCAGGCAGAG CAGGCCGGA AGGAGCAGGT TGGGAGTGTG GTGGGGCCCCG CAGCTTCAGC | 120 |



4

```

AGTGCCGCGG TGA CTATGGC CCCGATCAAG GTGGGAGACA CCATTCCTC AGTGGAGGTA      180
TTTGRAGGGG AACCTGGAAA GAAGGTGAAC TTGGCAGAGC TGTTCAGGA CAAGAAAGGT      240
GTTTTGTTTG GAGTCCCTGG GGCATTTACA CCTGGCTGTT CCAAGACCCA TCTGCCTGGG      300
TTTGTGGAGC AAGCCGGAGC TCYGAAGGCC AAGGGAGCAC AAGTGGTGGC CTGTCTGAGT      360
GTTAATGATG YCTTCGTGAC TGCAGAGTGG GGTGAGGCC ACCAGGCAGA AGGCAAGGTT      420
CAGCTCCTGG CTGACCCAC TGGAGCTTTT GGAAAGGAGA CAGATTTACT ACTAGATGAT      480
TCTTTGGTGT CTCTCTTTGG GAATCGTCGG CTAAAAAGGT TCTCCATGGT GATAGACAAG      540
GGCGTAGTAA AGGCACTGAA CGTGGAGCCG GATGGCACAG GCCTCACCTG CAGCCTGGCC      600
CCCAACATCC TCTCACAAC CTGAGGCCCT GACCAGAATG TCCTCTGACT CTCCCATCTC      660
CTCCACCCAG CTCTGGGCCA AAGGCCCAGT ACCTCCTTAC CTGAGGGCCA CTGGAATGGA      720
ACCTTGACAA TATTTCTGCA ATAAACAGTT TAATTTGTGA AAAAAAAAAA AAAAAAAAAA      780

```

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 162 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Rattus Rattus

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 17
- (D) OTHER INFORMATION: /product= "Glu or Gly"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 63
- (D) OTHER INFORMATION: /product= "Leu or Pro"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 79
- (D) OTHER INFORMATION: /product= "Ala or Val"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

```

Met Ala Pro Ile Lys Val Gly Asp Thr Ile Pro Ser Val Glu Val Phe
1           5           10           15

```



5

Xaa Gly Glu Pro Gly Lys Lys Val Asn Leu Ala Glu Leu Phe Lys Asp
 20 25 30
 Lys Lys Gly Val Leu Phe Gly Val Pro Gly Ala Phe Thr Pro Gly Cys
 35 40 45
 Ser Lys Thr His Leu Pro Gly Phe Val Glu Gln Ala Gly Ala Xaa Lys
 50 55 60
 Ala Lys Gly Ala Gln Val Val Ala Cys Leu Ser Val Asn Asp Xaa Phe
 65 70 75 80
 Val Thr Ala Glu Trp Gly Arg Ala His Gln Ala Glu Gly Lys Val Gln
 85 90 95
 Leu Leu Ala Asp Pro Thr Gly Ala Phe Gly Lys Glu Thr Asp Leu Leu
 100 105 110
 Leu Asp Asp Ser Leu Val Ser Leu Phe Gly Asn Arg Arg Leu Lys Arg
 115 120 125
 Phe Ser Met Val Ile Asp Lys Gly Val Val Lys Ala Leu Asn Val Glu
 130 135 140
 Pro Asp Gly Thr Gly Leu Thr Cys Ser Leu Ala Pro Asn Ile Leu Ser
 145 150 155 160
 Gln Leu

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 675 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Mouse

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 99..588

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

TGCTCCGTGC ATCGACGTGC TTGGCAGGCA GAGCAGGCCG GAAAGAAGCA GGTGGGAGT 60
 GTGGCGGAGC CCGCAGCTTC AGCAGCTCCG CGGTGACCAT GGCCCCGATC AAGGTGGGAG 120
 ATGCCATTCC CTCAGTGGAG GTATTTGAAG GGGAACCGGG AAAGAAGGTG AACTTGGCAG 180
 AGCTGTTCAA GGGCAAGAAA GGTGTTTTGT TTGGAGTCCC TGGGGCATTT ACACCTGGCT 240


```

GTTCTAAGAC CCACCTGCCT GGGTTTGTGG AGCAAGCTGG AGCTCTGAAG GCTAAGGGAG      300
CGCAGGTGGT GGCCTGTCTG AGCGTTAATG ACGTCTTTGT GATTGAAGAG TGGGGTCGAG      360
CCCACCAGGC AGAAGGCAAG GTTCGGCTCC TGGCTGACCC CACTGGAGCC TTTGGGAAGG      420
CGACAGACTT ATTATTGGAT GATTCTTTGG TGTCTCTCTT TGGGAATCGT CGGCTGAAAA      480
GGTTCTCCAT GGTGATAGAC AACGGCATAG TGAAGGCACT GAACGTGGAG CCAGATGGCA      540
CAGGCCTCAC CTGCAGCCTG GCCCCAACA TCCTCTCCCA ACTCTGAGGC CCTGGCCAGA      600
TGTCCTCTGA CTCTCCCATC TCTCCCACCC GGCTCTAGGC CAAAAGGCTC GGTACCTCCT      660
TACTGGGAGC CACGT                                          675

```

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 162 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Mouse

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

```

Met Ala Pro Ile Lys Val Gly Asp Ala Ile Pro Ser Val Glu Val Phe
1           5           10           15
Glu Gly Glu Pro Gly Lys Lys Val Asn Leu Ala Glu Leu Phe Lys Gly
20           25           30
Lys Lys Gly Val Leu Phe Gly Val Pro Gly Ala Phe Thr Pro Gly Cys
35           40           45
Ser Lys Thr His Leu Pro Gly Phe Val Glu Gln Ala Gly Ala Leu Lys
50           55           60
Ala Lys Gly Ala Gln Val Val Ala Cys Leu Ser Val Asn Asp Val Phe
65           70           75           80
Val Ile Glu Glu Trp Gly Arg Ala His Gln Ala Glu Gly Lys Val Arg
85           90           95
Leu Leu Ala Asp Pro Thr Gly Ala Phe Gly Lys Ala Thr Asp Leu Leu
100          105          110
Leu Asp Asp Ser Leu Val Ser Leu Phe Gly Asn Arg Arg Leu Lys Arg
115          120          125

```


Phe Ser Met Val Ile Asp Asn Gly Ile Val Lys Ala Leu Asn Val Glu
 130 135 140

Pro Asp Gly Thr Gly Leu Thr Cys Ser Leu Ala Pro Asn Ile Leu Ser
 145 150 155 160

Gln Leu

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 469 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 161..382

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

| | |
|--|-----|
| GGGTATGGGA CTAGCTGGCG TGTGCGCCCT GAGACGCTCA GCGGGCTATA TACTCGTCGG | 60 |
| TGGGGCCGGC GGTCAGTCTG CGGCAGCGGC AGCAAGACGG TGCAGTGAAG GAGAGTGGGC | 120 |
| GTCTGGCGGG GTCCGCAGTT TCAGCAGAGC CGCTGCAGCC ATGGCCCCAA TCAAGGTTCTG | 180 |
| GCTCCTGGCT GATCCCACTG GGGCCTTTGG GAAGGAGACA GACTTATTAC TAGATGATTC | 240 |
| GCTGGTGTCC ATCTTTGGGA ATCGACGTCT CAAGAGGTTC TCCATGGTGG TACAGGATGG | 300 |
| CATAGTGAAG GCCCTGAATG TGAACCAGA TGGCACAGGC CTCACCTGCA GCCTGGCACC | 360 |
| CAATATCATC TCACAGCTCT GAGGCCCTGG GCCAGATTAC TTCCTCCACC CCTCCCTATC | 420 |
| TCACCTGCCC AGCCGTGTGC TGGGGCCCTG CAATTGGAAT GTTGCCAG | 469 |

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 601 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION:161..514

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

| | |
|---|-----|
| GGGTATGGGA CTAGCTGGCG TGTGCGCCCT GAGACGCTCA GCGGGCTATA TACTCGTCGG | 60 |
| TGGGGCCGGC GGTCACTCTG CGGCAGCGGC AGCAAGACGG TGCAGTGAAG GAGAGTGGGC | 120 |
| GTCTGGCGGG GTCCGCAGTT TCAGCAGAGC CGCTGCAGCC ATGGCCCCAA TCAAGACACA | 180 |
| CCTGCCAGGG TTTGTGGAGC AGGCTGAGGC TCTGAAGGCC AAGGGAGTCC AGGTGGTGGC | 240 |
| CTGTCTGAGT GTTAATGATG CCTTTGTGAC TGGCGAGTGG GGCCGAGCCC ACAAGGCGGA | 300 |
| AGGCAAGGTT CGGCTCCTGG CTGATCCCAC TGGGGCCTTT GGAAGGAGA CAGACTTATT | 360 |
| ACTAGATGAT TCGCTGGTGT CCATCTTTGG GAATCGACGT CTCAAGAGGT TCTCCATGGT | 420 |
| GGTACAGGAT GGCATAGTGA AGGCCCTGAA TGTGGAACCA GATGGCACAG GCCTCACCTG | 480 |
| CAGCCTGGCA CCAATATCA TCTCACAGCT CTGAGGCCCT GGGCCAGATT ACTTCCTCCA | 540 |
| CCCCTCCCTA TCTCACCTGC CCAGCCCTGT GCTGGGGCCC TGCAATTGGA ATGTTGGCCA | 600 |
| G | 601 |

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 604 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION:161..517

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

| | |
|---|----|
| GGGTATGGGA CTAGCTGGCG TGTGCGCCCT GAGACGCTCA GCGGGCTATA TACTCGTCGG | 60 |
|---|----|

| | | | | | | |
|------------|------------|------------|------------|------------|------------|-----|
| TGGGGCCGGC | GGTCAGTCTG | CGGCAGCGGC | AGCAAGACGG | TGCAGTGAAG | GAGAGTGGGC | 120 |
| GTCTGGCGGG | GTCCGCAGTT | TCAGCAGAGC | CGCTGCAGCC | ATGGCCCCAA | TCAAGGTGGG | 180 |
| AGATGCCATC | CCAGCAGTGG | AGGTGTTTGA | AGGGGAGCCA | GGGAACAAGG | TGAACCTGGC | 240 |
| AGAGCTGTTT | AAGGGCAAGA | AGGGTGTGCT | GTTTGGAGTT | CCTGGGGCCT | TCACCCCTGG | 300 |
| ATGTTCCAAG | GTTCGGCTCC | TGGCTGATCC | CACTGGGGCC | TTTGGGAAGG | AGACAGACTT | 360 |
| ATTACTAGAT | GATTCGCTGG | TGTCCATCTT | TGGGAATCGA | CGTCTCAAGA | GGTTCTCCAT | 420 |
| GGTGGTACAG | GATGGCATAG | TGAAGGCCCT | GAATGTGGAA | CCAGATGGCA | CAGGCCTCAC | 480 |
| CTGCAGCCTG | GCAQCCAATA | TCATCTCACA | GCTCTGAGGC | CCTGGGCCAG | ATTACTTCCT | 540 |
| CCACCCCTCC | CTATCTCACC | TGCCCAGCCC | TGTGCTGGGG | CCCTGCAATT | GGAATGTTGG | 600 |
| CCAG | | | | | | 604 |

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2710 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(ix) FEATURE:

- (A) NAME/KEY: exon
- (B) LOCATION: 2516..2710

(ix) FEATURE:

- (A) NAME/KEY: exon
- (B) LOCATION: 2074..2135

(ix) FEATURE:

- (A) NAME/KEY: exon
- (B) LOCATION: 1932..1970

(ix) FEATURE:

- (A) NAME/KEY: exon
- (B) LOCATION: 1728..1859

(ix) FEATURE:

- (A) NAME/KEY: exon
- (B) LOCATION: 802..936

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

| | | | | | | |
|------------|------------|------------|------------|------------|------------|------|
| TCTGTCCCTT | AGCGCCCCCG | CGGGGGCTTA | CCCCATCCCA | CTCCATGACC | TCCCCTCCCC | 60 |
| CCATGGCGAA | TTCCCACCTT | TCTGTCTTTC | ACTCACTTCC | TGGAACCGTC | CCCAGGGCCT | 120 |
| TGGACCTTCC | CCCTTCTCCT | CCCAAACCTT | GTGAGACCCC | ATTCCCTTTC | TACTTCATCC | 180 |
| TGCTCTCAAC | TTTTGGGCTC | CTCAGAGGCC | CTCACCCTTG | ACTCTCTCTC | CCTACCCACT | 240 |
| CTGGTCCCAT | GAAGCCCTCA | AGTACTCTGG | GGATGGATCC | TTCCCCCTTC | AAAAGATTCC | 300 |
| TTCTTTTGTT | CTACACCTCC | TGGGTGTAGG | GGCCTGGACA | CCCTCCCCCA | ACGTTCCACC | 360 |
| TGCCGCTGCC | CTTCCTCTTC | CTCCTCCTGA | GGGTGGGACC | CTCAGACCTG | GCCAAGATCC | 420 |
| TCTCCCTCCA | TGTTGTCAGG | GACTCCTCCT | CACCCCCAAA | TACAGCCCTC | TAGCCCCTGT | 480 |
| CCATTTTATT | CCACTCCTTT | CCTGTAACCT | AGACAGCATG | TTATGCAACC | CTTTGCGACA | 540 |
| CATGGGGAAA | CCTTCCCTCC | CTTCCTCTGT | TGTCACCAAT | GGCCCCTTAA | GAGGAGCAGG | 600 |
| GCCACCTTGA | AACTTGAGG | ATATGGGGTA | ACCCAGTGGG | AGCGGGCAGG | GAGGGCCCTT | 660 |
| GGAAACTGAC | AGGGCTGGAG | TATCCTGCTG | GGTTTCAGCC | CCGGTTCCTG | CAGGCACAGC | 720 |
| TGCCAGGCTC | TCTGTTCAAC | TTCCTGCCTC | TGGTTTGCCC | CGGCTCCCTC | ACCCCCCTTA | 780 |
| CCCTGGAGTC | CTTCCTTCTA | GGTGGGAGAT | GCCATCCCAG | CAGTGGAGGT | GTTTGAAGGG | 840 |
| GAGCCAGGGA | ACAAGGTGAA | CCTGGCAGAG | CTGTTCAAGG | GCAAGAAGGG | TGTGCTGTTT | 900 |
| GGAGTTCCTG | GGGCCTTCAC | CCCTGGATGT | TCCAAGGTGA | GGCCCTTCCC | CTTCTGAAGA | 960 |
| TCAGGACCTG | GGGATCTTTT | GTGTTGCTCT | TAAGTCCTCC | ACATAGTCCT | GATAGGACTC | 1020 |
| CTAAAAAGCA | TTTCAGTGCC | ATCACAAAAC | AAGTAGAGCT | GGGTAGAGCT | GGGCGCGGTG | 1080 |
| GCTCACGCCT | GTAATCCCAG | CACCTTGGGA | GGCCAAGGCG | GGTGGATCAC | GAGGTCAGGA | 1140 |
| GTCCAAAACC | AGCCTGGCCA | AGATGGTGAA | ACCCTGTCTC | TACTAAAAAT | GCAAAAAAAT | 1200 |
| CAGCCGGATA | TGGTGGCGGG | CGCCTGTAAT | CCCAGGTATT | GGGGAGGCTG | AGGCAGAGAA | 1260 |
| TTGCTTGAAC | CCAGGAGGCG | TAGGTTGCAG | TGAGTGGAGA | TCGTGCCTCT | GCAGTCCAGC | 1320 |
| CTGGGTGAAA | GAGCGAGACT | CCGTCTCAAA | ATGAAAAAAA | AAAAAGAAAA | CAAGTAGAGA | 1380 |
| CTGCAAAAAG | GGAACAGTAC | CGGGAATGTT | GGAGAAAAAC | ATACTACAAT | TAAATCCAAC | 1440 |
| ACCCCTGTTG | GTCCTGCTAA | ATGACAGGCA | CTGTGGAAGG | TGCTTGGGAC | TCAGATAAAT | 1500 |
| AAGACAAAGA | TCTGCCCATG | GAAAGTTCAC | GTCTGGACCA | TAAGGCATTA | GGTTTCATTC | 1560 |
| TGAGCTTCCT | AGTGGCCAAG | GCAAAAAGGA | AATAGAATGG | TTTAGACAGC | TCTATTGTC | 1620 |
| TGATCAAAGG | TGTTGAGGCA | GAGCACTGAG | GAGGGCCTGG | AGATAAAGGG | TGGGCTGGGG | 1680 |
| GTCAGATGCA | GTTATCCCTT | TGCCGACCCT | TTGTTCCCCT | TCCTCAGACA | CACCTGCCAG | 1740 |
| GGTTTGTGGA | GCAGGCTGAG | GCTCTGAAGG | CCAAGGGAGT | CCAGGTGGTG | GCCTGTCTGA | 1800 |
| GTGTTAATGA | TGCCTTTGTG | ACTGGCGAGT | GGGGCCGAGC | CCACAAGGCG | GAAGGCAAGG | 1860 |

| | |
|---|------|
| TGAGGTGAGG GGCCTGCAGG GAGTCAGGAC CAGGTAGGAT ATTCTTCTTG TGACCTCTAC | 1920 |
| TTTCTCTGCA GGTTCGGCTC CTGGCTGATC CCACTGGGGC CTTTGGGAAG GTGAGTGTTC | 1980 |
| CCCTGACCGC CACAGGGACA TGGCGGTGCG GGGAGCAGTG GGGGCCCTTG GCCTCTTCAA | 2040 |
| GGATTTCTGA CACTTTTCTC TGTCTCTTCT TAGGAGACAG ACTTATTACT AGATGATTCG | 2100 |
| CTGGTGTCCA TCTTTGGGAA TCGACGTCTC AAGAGGTAAA AGTGGAGAGT CCTCTGTGGA | 2160 |
| GAAAGTCCTC TGTGGGAGAG AGTCCTCTGT GGGAGAGAGT CCTCTGTGGA GAGGGTCCTC | 2220 |
| TGTGGGAAGA GTCGTCTGTG GGGGAGATGT GTGGGAGAGA GTCCTGTGTG GGGAGAGTCT | 2280 |
| TCTGTAGGGG AGAGTCCTCT GGGGAGAGAG TCCTGTGTGG GGGAGAGTCC TCTGTGGGGA | 2340 |
| GAGTCCTCTG TGTGGAGAGA GTCCTGTGTG GTGGTGAGTC CTCTGTGGGG GAGAGTCCTC | 2400 |
| TGTGGGGGGA GTCCTCTCTG GAGTTCTCTT GGGCCCCTGG CTGTTCACTG CCTGTCTCCA | 2460 |
| TGCCCAGCCT CCAAGCCCAG GCTGATGCAG CTGGCTGGGC CCCTCTTTCC GGCAGGTTCT | 2520 |
| CCATGGTGGT ACAGGATGGC ATAGTGAAGG CCCTGAATGT GGAACCAGAT GGCACAGGCC | 2580 |
| TCACCTGCAG CCTGGCACCC AATATCATCT CACAGCTCTG AGGCCCTGGG CCAGATTACT | 2640 |
| TCCTCCACCC CTCCCTATCT CACCTGCCCA GCCCTGTGCT GGGGCCCTGC AATTGGAATG | 2700 |
| TTGGCCAGAT | 2710 |

(2) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

GCCATCCCAG CAGTGGAGGT GTTTG

25

(2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

12

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

TTGAACAGCT CTGCCAGGTT CACC

24

(2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

TGGAGGTGTT TGAAGGGGAG CCAG

24

(2) INFORMATION FOR SEQ ID NO: 14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

CAGGTTTACC TTGTTCCCTG GCTC

24

(2) INFORMATION FOR SEQ ID NO: 15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

GGGTATGGGA CTAGCTGGCG

20

(2) INFORMATION FOR SEQ ID NO: 16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single

13

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

CTGGCCAACA TTCCAATTGC AG

22

(2) INFORMATION FOR SEQ ID NO: 17:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

ATGTTATGCA ACCCTTTGCG ACAC

24

(2) INFORMATION FOR SEQ ID NO: 18:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

GTGTTTGAAG GGGAGCCAGG GAAC

24

(2) INFORMATION FOR SEQ ID NO: 19:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

AGAGACAGGG TTTCACCATC TTGG

24

1/7




FIG. 1

430 Rec'd [REDACTED] T/PTO 22 FEB 2000

2/7





FIG. 2


430 Rec'd [redacted] T/PTO 22 FEB 2000

3/7




430 Rec'd CT/PTO 22 FEB 2000

Gene (chromosome 11q12-13)



mRNAs



Proteins




FIG. 4

430 Rec'd CT/PTO 2 2 FEB 2000

CLUSTAL V alignment of human and rat B18 amino acid sequences (Identity: 90%, Homology: 97.5%):

```
B10hum      MAPIKVGDALPAVEVFEGEPGNKVNLAELFKGKGGVLFVPGAFTPGCSK = SEQIDNO1
B10rat      MAPIKVGDTIPSVEVFEGEPGNKVNLAELFKDKKGGVLFVPGAFTPGCSK
*****
```

B18hum THLPGFVEQAEALKAKGVQVVACLSVNDAFVTGEWGRAHKAEGKVRLLD
B18rat THLPGFVEQAGALKAKGAQVVACLSVNDVFVTAEWGRAHQAEGKVQLLD

FIG. 5a

```

B18hum      PTGAFGKETDLLLDDSLVSI FGNRRLKRFSMVVDGIVKALNVEPDGTGL
B18rat      PTGAFGKETDLLLDDSLVSLFGNRRLKRFSMVIDKGVVKALNVEPDGTGL
*****

```

```

B18hum      TCSLAPNIIISQL
B18rat      TCSLAPNIIISQL
            *****

```

CLUSTAL V alignment of human and mouse B18 amino acid sequences (Identity: 91%, Homology: 96%):

```

B18hum      MAPIKVGDAIPAVEVFEGEPGNKVNLAELFKGKKGVLFVGPGAFTPGCSK
B18mouse    MAPIKVGDAIPSEVFEGEPGKKVNLAELFKGKKGVLFVGPGAFTPGCSK
*****

```

```
B18hum      THLPGFVEQAEALKAKGVQVVAACLSVNDAFVTGEWGRAHKAEGKVRLLAD
B18mouse    THLPGFVEQAGALKAKGAQVVAACLSVNDVFVI EEWGRAHQAEKGVRLLAD
            *-----*
```

```

B18hum      PTGAFGKETDLLLDDSLVSI FGNRRRLKRFSMVVDGIVKALNVEPDGTGL
B18mouse    PTGAFGKATDLLLDDSLVSLFGNRRRLKRFSMVIDNGIVKALNVEPDGTGL
*****

```

```

B18hum      TCSLAPNIIISQL
B18mouse    TCSLAPNIIISQL
            *****

```

CLUSTAL V alignment of human and rat cDNA sequences (identity: 612/780, 78.5%):

B18hum GCCAGGAGGCGGAGTGGAAAGTGGCCGTGGGGCGGGTATGGGACTAGCTGG
B18rat -----TG-----CGTC-----CTAGGCAG

B18hum CGTGTGCGCCCTGAGACGCTCAGCGGGCTATATACTCGTCGGTG GGGGCCG
B18rat CATA---GCC---GGA---TCGGTGCTCCGTGCATCGGCTACTTGGAC--
* * * * * * * * * *

```

B18hum      GCGGTCAGTCTGCGGCAGCGGCAGCAAGACGGTGCACTGAAGGAGAGTGG
B18rat      -----GTGCGTGGCAGGCAGAGCAGGCCCGG---AAAGGAGCAGGTTGG
              * * * * *      * * * * *      * * * * *      * * * * *

```

430 Rec'd PQT/PTO 2 2 FEB 2000

6/7

FIG.5b

B18hum GCGTCTGGCGGGGTCCGCAGTTTCAGCAGAGCCGCTGCAGCCATGGCCCC
B18rat GAGTGTGGTGGGGCCCGCAGCTTCAGCAGTGCCGCGGTGACTATGGCCCC
* * * * *

B18hum AATCAAGGTGGGAGATGCCATCCCAGCAGTGGAGGTGTTTGAAGGGGAGC
B18rat GATCAAGGTGGGAGACACCATTCCCTCAGTGGAGGTATTTGAAGGGGAAC
* * * * *

B18hum CAGGGAACAAGGTGAACCTGGCAGAGCTGTTCAAGGGCAAGAAGGGTGTG
B18rat CTGGAAGAAGGTGAACTTGGCAGAGCTGTTCAAGGACAAGAAAGGTGTT
* * * * *

B18hum CTGTTTGGAGTTCCTGGGGCCTTCACCCCTGGATGTTCCAAGACACACCT
B18rat TTGTTTGGAGTCCCTGGGGCATTACACCTGGCTGTTCCAAGACCCATCT
* * * * *

B18hum GCCAGGGTTTGTGGAGCAGGCTGAGGCTCTGAAGGCCAAGGGAGTCCAGG
B18rat GCCTGGGTTTGTGGAGCAAGCCGGAGCTCTGAAGGCCAAGGGAGCACAAAG
* * * * *

B18hum TGGTGGCCTGTCTGAGTGTTAATGATGCCTTTGTGACTGGCGAGTGGGGC
B18rat TGGTGGCCTGTCTGAGTGTTAATGATGTCTTCGTGACTGCAGAGTGGGGT
* * * * *

B18hum CGAGCCCACAAGGCGGAAGGCAAGGTTCCGGCTCCTGGCTGATCCCCTGAG
B18rat CGAGCCCACCAGGCAGAAGGCAAGGTTCCAGCTCCTGGCTGACCCCTGAG
* * * * *

B18hum GGCCTTTGGGAAGGAGACAGACTTATTACTAGATGATTGCTGGTGTCCA
B18rat AGCTTTTGGAAAGGAGACAGATTACTACTAGATGATTCTTTGGTGTCTC
* * * * *

B18hum TCTTTGGGAATCGACGTCTCAAGAGGTTCTCCATGGTGGTACAGGATGGC
B18rat TCTTTGGGAATCGTCGGCTAAAAAGGTTCTCCATGGTGATAGACAAGGGC
* * * * *

B18hum ATAGTGAAGGCCCTGAATGTGGAACCAGATGGCACAGGCCTCACCTGCAG
B18rat GTAGTAAAGGCACTGAACGTGGAGCCGGATGGCACAGGCCTCACCTGCAG
* * * * *

B18hum CCTGGCACCCAATATCATCTCACAGCTCTGAGGCCCTGGGCCAGATTACT
B18rat CCTGGCCCCCAACATCCTCTCACAACCTCTGAGGCCCTGA-CCAGA--ATG
* * * * *

B18hum TCCTCCACCCCTCCCTATCTCACCTGCCAGCCCTGTGCTGG-GGCCCTG
B18rat TCCTCTGACTCTCCC-ATCTCCTCCACCCAGCTCTGGGCCAAAGGCCAG
* * * * *

B18hum CA-----ATTGGAATG-----TTGGCCAGATTTCTGC
B18rat TACCTCCTTACCTGAGGGCCACTGGAATGGAACCTTGACAATATTTCTGC
* * * * *

B18hum AATAAACACTTGTGGTTTGC GGAAAAA-----
B18rat AATAAACAGTT-TAATTTGTGAAAAAAAAAAAAAAAAAAAA
* * * * *

430 Rec'd PCT/PTO 2 2 FEB 2000

7/7

CLUSTAL V alignment of human and mouse cDNA sequences (Identity: 552/675, 81.8%):

FIG.5c

```
B18hum      GCCAGGAGGCGGAGTGGAAAGTGGCCGTGGGGCGGGTATGGGACTAGCTGG
B18mouse    -----

B18hum      CGTGTGCGCCCTGAGACGCTCAGCGGGCTATATACTCGTCGGTGGGGCCG
B18mouse    -----TGCTCCGTG-----CATCGACGTGCTTG
                  **** *

B18hum      GCGGTCAGTCTGCGGCAGCGGCAGCAAGACGGTGCAGTGAAGGAGAGTGG
B18mouse    GCAGGCAG-----AGCAGGCCGG---AAAGAAGCAGGTTGG
                  ** * **** * **** *

B18hum      GCGTCTGGCGGGGTCCGCAGTTTCAGCAGAGCCGCTGCAGCCATGGCCCC
B18mouse    GAGTGTGGCGGAGCCCGCAGCTTCAGCAGCTCCGCGGTGACCATGGCCCC
                  * ** *

B18hum      AATCAAGGTGGGAGATGCCATCCCAGCAGTGGAGGTGTTTGAAGGGGAGC
B18mouse    GATCAAGGTGGGAGATGCCATTCCCTCAGTGGAGGTATTTGAAGGGGAAC
                  *****

B18hum      CAGGGAACAAGGTGAACCTGGCAGAGCTGTTCAAGGGCAAGAAGGGTGTG
B18mouse    CGGGAAAGAAGGTGAACCTGGCAGAGCTGTTCAAGGGCAAGAAAGGTGTT
                  * * *

B18hum      CTGTTTGGAGTTCCTGGGGCCTTCACCCCTGGATGTTCCAAGACACACCT
B18mouse    TTGTTTGGAGTCCCTGGGGCATTACACCTGGCTGTTCTAAGACCCACCT
                  *****

B18hum      GCCAGGGTTTGTGGAGCAGGCTGAGGCTCTGAAGGCCAAGGGAGTCCAGG
B18mouse    GCCTGGGTTTGTGGAGCAAGCTGGAGCTCTGAAGGCTAAGGGAGCGCAGG
                  ***

B18hum      TGGTGGCCTGTCTGAGTGTAAATGATGCCTTTGTGACTGGCGAGTGGGGC
B18mouse    TGGTGGCCTGTCTGAGCGTTAATGACGTCTTTGTGATTGAAGAGTGGGGT
                  *****

B18hum      CGAGCCCACAAGGCGGAAGGCAAGGTTCCGGCTCCTGGCTGATCCCACTGG
B18mouse    CGAGCCCACCAGGCAGAAGGCAAGGTTCCGGCTCCTGGCTGACCCCACTGG
                  *****

B18hum      GGCCTTTGGGAAGGAGACAGACTTATTACTAGATGATTCGCTGGTGTCCA
B18mouse    AGCCTTTGGGAAGGCGACAGACTTATTATTGGATGATTCTTTGGTGTCTC
                  *****

B18hum      TCTTTGGGAATCGACGTCTCAAGAGGTTCTCCATGGTGGTACAGGATGGC
B18mouse    TCTTTGGGAATCGTCGGCTGAAAAGGTTCTCCATGGTGATAGACAACGGC
                  *****

B18hum      ATAGTGAAGGCCCTGAATGTGGAACCAGATGGCACAGGCCTCACCTGCAG
B18mouse    ATAGTGAAGGCACTGAACGTGGAGCCAGATGGCACAGGCCTCACCTGCAG
                  *****

B18hum      CCTGGCACCCAATATCATCTCACAGCTCTGAGGCCCTGGGCCAGATTACT
B18mouse    CCTGGCCCCCAACATCCTCTCCCAACTCTGAGGCCCTGG-CCAGATG---
                  *****

B18hum      TCCTCCACCCCTCCCTATCTCACCTGCCAGCCCTGTGCTGGGGCCCTGC
B18mouse    TCCTCTGACTCTCC-ATCTCTCCACCCGGCTCT-----AGGCC-----
                  *****


B18hum      AATTGGAATGTTGGCCAGATTTCTGCAATAAACACTTGTGGTTTGGCGAA
B18mouse    ----AAAAGGCTCGGTACCTCCTTACTGGGAGC-CACGT-----
                  *****
```




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)


| | | | |
|--|--|---|--|
| (51) International Patent Classification ⁶ : C07K 14/00 | | A2 | (11) International Publication Number: WO 99/09054 |
| | | | (43) International Publication Date: 25 February 1999 (25.02.99) |
| (21) International Application Number: PCT/BE98/00124 | | (81) Designated States: CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). | |
| (22) International Filing Date: 20 August 1998 (20.08.98) | | | |
| (30) Priority Data: 9700692 20 August 1997 (20.08.97) BE | | Published Without international search report and to be republished upon receipt of that report. | |
| (71) Applicants (for all designated States except US): UNIVERSITE CATHOLIQUE DE LOUVAIN [BE/BE]; Halles Universitaires, Place de l'Université 1, B-1348 Louvain-la-Neuve (BE). UNIVERSITE DE MONS-HAINAUT [BE/BE]; Place du Parc 20, B-7000 Mons (BE). | | | |
| (72) Inventors; and (75) Inventors/Applicants (for US only): KNOOPS, Bernard [BE/BE]; Rue Chapelle Notre-Dane 3/1, B-1341 Ceroux-Mousty (BE). HERMANS, Cedric [BE/BE]; Avenue des Glycines 42, B-1030 Brussels (BE). BERNARD, Alfred [BE/BE]; Avenue de la Chapelle 6, B-1200 Brussels (BE). WATTIEZ, Ruddy [BE/BE]; Chemin du Sauvelon 17, B-7022 Hyon (BE). FALMAGNE, Paul [BE/BE]; Rue du Point du Jour 8, B-7022 Mesvin (BE). | | | |
| (74) Agents: VAN MALDEREN, Eric et al.; Office Van Malderen, Place Reine Fabiola 6/1, B-1083 Brussels (BE). | | | |

(54) Title: PEROXISOME-ASSOCIATED POLYPEPTIDE, NUCLEOTIDE SEQUENCE ENCODING SAID POLYPEPTIDE AND THEIR USES IN THE DIAGNOSIS AND/OR THE TREATMENT OF LUNG INJURIES AND DISEASES, AND OF OXIDATIVE STRESS-RELATED DISORDERS



(57) Abstract

The present invention is related to an isolated and purified polypeptide which amino acid sequence presents more than 70 % with the sequence SEQ ID NO 1. The present invention is also related to the nucleotide sequence encoding said amino acid sequence, the inhibitor directed against said sequences and their use in the diagnosis, treatment and/or prevention of lung injuries or diseases and oxidative stress-related disorders.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | ML | Mali | TR | Turkey |
| BG | Bulgaria | HU | Hungary | MN | Mongolia | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MR | Mauritania | UA | Ukraine |
| BR | Brazil | IL | Israel | MW | Malawi | UG | Uganda |
| BY | Belarus | IS | Iceland | MX | Mexico | US | United States of America |
| CA | Canada | IT | Italy | NE | Niger | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NL | Netherlands | VN | Viet Nam |
| CG | Congo | KE | Kenya | NO | Norway | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NZ | New Zealand | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | PL | Poland | | |
| CM | Cameroon | KR | Republic of Korea | PT | Portugal | | |
| CN | China | KZ | Kazakhstan | RO | Romania | | |
| CU | Cuba | LC | Saint Lucia | RU | Russian Federation | | |
| CZ | Czech Republic | LI | Liechtenstein | SD | Sudan | | |
| DE | Germany | LK | Sri Lanka | SE | Sweden | | |
| DK | Denmark | LR | Liberia | SG | Singapore | | |
| EE | Estonia | | | | | | |

5

10 PEROXISOME-ASSOCIATED POLYPEPTIDE, NUCLEOTIDE SEQUENCE
ENCODING SAID POLYPEPTIDE AND THEIR USES IN THE DIAGNOSIS
AND/OR THE TREATMENT OF LUNG INJURIES AND DISEASES, AND OF
OXIDATIVE STRESS-RELATED DISORDERS

Field of the invention

15 The present invention is related to a new peroxisome-associated polypeptide, the nucleotide sequence encoding said polypeptide and portions thereof as well as their uses in the diagnosis of several diseases, especially the diagnosis and/or the treatment of lung injuries and
20 diseases, and of oxidative stress-related disorders.

Background of the invention

The peroxisomes are organelles nearly ubiquitous in eukaryotic cells. They contain enzymes
25 essential for various catabolic and anabolic pathways. Some of these enzymes are expressed constitutively while others can be induced under appropriate conditions. Peroxisomes carry out a variety of essential reactions such as peroxisomal oxidation and respiration, fatty acid beta-
30 oxidation, cholesterol and dolichol metabolism, ether-phospholipid synthesis, and glyoxylate and pipecolic acid metabolism.

The peroxisomal respiratory pathway is based upon the formation of hydrogen peroxide by a collection of oxidases and the decomposition of the H_2O_2 by catalase. These reactions are responsible for 20% of oxygen consumption in liver, and several oxidases have been identified in peroxisomes. Ethanol elimination via catalase in peroxisomes may be significant in addition to the oxidation via cytosolic alcohol dehydrogenase.

The peroxisomal beta-oxidation system catalyses the beta-oxidative chain shortening of a specific set of compounds which can not be handled by mitochondria : very long chain fatty acids, di- and trihydroxycholestanoic acids, pristanic acid, long chain dicarboxylic acids, several prostaglandins, several leukotrienes, 12- and 15-hydroxyeicosatetraenoic acid, and several mono- and polyunsaturated fatty acids, which are of direct diagnostic relevance for some peroxisomal disorders.

Peroxisomes play also a major role in the synthesis of cholesterol and other isoprenoids. Fibroblasts from patients affected by disorders of peroxisome biogenesis show low capacity to synthesise cholesterol.

Two enzyme activities responsible for introduction of the characteristic ether linkage in ether-linked phospholipids (dihydroacetonephosphate acyltransferase (DHAPAT) and alkyl-dihydroxyacetonephosphate synthase (alkyl-DHAP synthase)) are localised in peroxisomes. These enzymes are not yet cloned. As demonstrated by the identification of patients with deficiency of either DHAPAT or alkyl-DHAP synthase with severe clinical abnormalities, ether-phospholipids are of major importance in humans.


Peroxisomes are able to detoxify glyoxylate via alanine/glyoxylate aminotransferase. The deficiency of this cloned enzyme causes hyperoxaluria type I.

L-pipecolate is a minor metabolite of L-lysine and is
5 catabolised by the L-pipecolate oxidase localised in peroxisomes. The enzyme is deficient in cerebro-hepato-renal (Zellweger) syndrome.

In human, the importance of peroxisomes was emphasised by a number of inherited diseases involving
10 either a defect in the biogenesis of peroxisomes or a deficiency of one (or more) peroxisomal enzymes. So far, 12 different peroxisomal disorders have been described and most of them are lethal.

A wide variety of chemicals have been showed
15 to produce peroxisome proliferation and induction of peroxisomal and microsomal fatty acids-oxidising enzymes activities in rats and mice. Several peroxisomes proliferators have been shown to increase the incidence of liver tumours in these species. Proposed mechanisms of
20 liver tumour formation by peroxisomes proliferators include induction of sustained oxidative stress.

Therefore, newly identified molecules associated with peroxisomes could be used for the development of diagnostic tools and possibly for the
25 improvement of several therapeutical applications of various diseases associated with peroxisomal disorders. In addition, it is useful to identify the molecules present in specific organs like the lung and which may be used as specific markers of inflammatory diseases as well as lung
30 injuries or diseases.



Summary of the invention

The Inventors have isolated and purified a new sequence of a low molecular weight human broncho-alveolar polypeptide. Said mammal, preferably human, protein or polypeptide (hereafter identified as B18hum
5 protein) has been sequenced and its corresponding genomic DNA (SEQ ID NO 8) and cDNA (SEQ ID NO 1) have been identified. Similarly, the corresponding nucleotide and amino acid sequence from a rat (SEQ ID NO 3 and 4) and from
10 a mouse (SEQ ID NO 5 and 6) have been obtained.

Said sequences present several homologies with other peroxisomal proteins of yeast and comprise a carboxy-terminal tripeptide SQL which is necessary for the specific targeting and translocation of several proteins
15 into the peroxisome.

Therefore, the present invention is related to a new isolated and purified polypeptide sequence having a amino acid sequence which presents more than 70% homology, advantageously more than 85% homology, more
20 preferably more than 95% homology, with the amino acid sequence SEQ ID NO 2. Said amino acid sequence is advantageously obtained from a mammal, preferably from a rat, a mouse or a human.

The present invention is also related to the
25 isolated and purified polypeptide sequence corresponding to the amino acid sequence SEQ ID NO 2 or a portion thereof, preferably an immunoreactive portion (putative immunogenic domain or T or B cell epitopes).

Said portions are advantageously comprised
30 between :

- Glutamic acid position 13 - Glutamic acid position 27
- Alanine position 26 - Leucine position 36

- Alanine position 42 - Glutamic acid position 57
- Glutamic acid position 57 - Valine position 69
- Valine position 80 - Leucine position 97
- Arginine position 95 - Leucine position 112
- 5 - Serine position 118 - Serine position 129
- Valine position 137 - Threonine position 150

Preferably, said portion has more than 10, 20, 30, 50 or 70 amino acids. Specific portions of the amino acid sequence SEQ ID NO 2 are also portions of more
10 than 70 amino acids which present at least 80% of the proteinic activity (see example 5) of the complete SEQ ID NO 2 sequence. Therefore, the amino acid sequence according to the invention can be partially deleted while maintaining its activity, preferably its anti-oxidative activity, which
15 will be described hereafter.

According to the invention, the amino acid sequence SEQ ID NO 2 presents a pI of 7.16 and a molecular weight of 17047 Dalton as hereafter defined by bidimensional electrophoresis.

20 The present invention is also related to the nucleotide sequence encoding the amino acid sequence according to the invention and its regulatory sequences upstream said coding sequence. A nucleotide sequence encoding the polypeptide according to the invention is a
25 genomic DNA (see SEQ ID NO 10), a cDNA (see SEQ ID NO 1) or a mRNA, possibly comprising said upstream regulatory sequence. Advantageously, said nucleotide sequence presents more than 70%, advantageously more than 85%, more preferably more than 95% homology with SEQ ID NO 1 or its
30 complementary strand.

According to a preferred embodiment of the present invention, said nucleotide sequence corresponds to the nucleotide sequence SEQ ID NO 1, its complementary strand or a portion thereof.

5 "A portion of the nucleotide sequence SEQ ID NO 1" means any nucleotide sequence of more than 15 base pairs (such as a primer, a probe or an antisense nucleotide sequence) which allow the specific identification, reconstitution or blocking of the complete nucleotide
10 sequence SEQ ID NO 1 (including its regulatory sequences upstream the coding sequence).

Said portions allow the specific identification, reconstitution or blocking by specific hybridisation with the nucleotidic sequence SEQ ID NO 1,
15 preferably under standard stringent conditions, with sequences like probes or primers possibly labelled with a compound (radioactive compound, enzyme, fluorescent marker, etc.), and can be used in a specific diagnostic or dosage method like probe hybridisation (see Sambrook et al., §§
20 9.47-9.51 in *Molecular Cloning : A Laboratory Manual*, Cold Spring Harbor, Laboratory Press, Cold Spring Harbor, New York (1989)), genetic amplification (like PCR (US patent 4,683,195), LCR (Wu et al., *Genomics* 4, pp. 560-569), CPR (US patent 5,011,769)).

25 Exemplary stringent hybridisation conditions are as follows : hybridisation at 42 °C in 50% formamide 5x SSC, 20 mM sodium phosphate, pH 6.8 washing in 0.2x SSC at 55 °C. It is understood by those skilled in the art that variation of these conditions occur based on the length and
30 GC nucleotide content of the sequence to be hybridised. Formulas standard in the art are appropriated for

determining exact hybridisation conditions (see Sambrook et al.

Preferred examples of said nucleotide portions are as follows :

| | <u>Sequence</u> | <u>Position</u> |
|----|---------------------------------|------------------------|
| 5 | 5'-gccatcccagcagtgagggtgtttg-3' | (SEQ ID NO 11) 217-241 |
| | 5'-ttgaacagctctgccaggttcacc-3' | (SEQ ID NO 12) 261-284 |
| | 5'-tgagggtgtttgaaggggagccag-3' | (SEQ ID NO 13) 230-253 |
| | 5'-caggttcaccttgttccctggctc-3' | (SEQ ID NO 14) 247-270 |
| 10 | 5'-gggtatgggactagctggcg-3' | (SEQ ID NO 15) 33-52 |
| | 5'-ctggccaacattccaattgcag-3' | (SEQ ID NO 16) 747-768 |

and the sequences of respectively 601 (SEQ ID NO 8), 604 (SEQ ID NO 9) and 469 (SEQ ID NO 7) base pairs corresponding to specific mRNA alternative splicing of the B18 human nucleotide sequence as described in Figure 4 (the known genomic sequence incorporating several introns and exons is represented in the sequence SEQ ID NO 10).

Said sequences may be used for a genetic amplification or a probe hybridisation as above-described.

The present invention is also related to a vector comprising the necessary elements for the injection, transfection or transduction of cells and having incorporated one or more of the nucleotide sequences according to the invention. The vector according to the invention is selected from the group consisting of viruses, plasmids, phagemides, cationic vesicles, liposomes or a mixture thereof. Said vector may comprise also one or more adjacent regulatory sequences (such as promoter(s), secretion and termination signal sequence(s)), advantageously operably linked to the nucleotide sequence according to the invention.

The present invention is also related to the cell transformed by said vector and expressing the polypeptide according to the invention.


The nucleotide sequence according to the invention can be also introduced in said cell by the formation of CaPO_4 -nucleic acid precipitate, DEAE-dextran-nucleic acid complex or by electroporation.

Another aspect of the present invention is related to an inhibitor of the polypeptide according to the invention or the nucleotide sequence according to the invention (including the upstream sequences like promoter-operator regulatory sequence which may be inhibited by a cis- and/or transactivating repressor). Said inhibitor is advantageously an antibody or a fragment of said antibody such as an hypervariable portion of said antibody directed against the amino acid or nucleotide sequence of the polypeptide according to the invention. Other examples of inhibitors according to the invention are antisense nucleotide sequences which allow the blocking of the expression of the nucleotide sequence according to the invention.

Another aspect of the present invention is related to a diagnostic device (such as a diagnostic kit or a chromatographic column) comprising an element selected from the group consisting of the amino acid sequence of said polypeptide, its nucleotide sequence, and/or the inhibitor according to the invention or a fragment thereof as above-described. Said diagnostic device may comprise also necessary reactants and media for the diagnostic and/or dosage of the nucleotide and/or amino acid sequence of the polypeptide according to the invention, which are based upon the method selected from the group consisting of

in situ hybridisation, hybridisation by labelled antibodies, especially RIA (Radio Immuno Assay) or ELISA (Enzymes Linked Immuno-Sorbent Assay) technologies, detection upon filter, upon solid support, in solution, in sandwich, upon gel, dot blot hybridisation, Northern blot hybridisation, Southern blot hybridisation, isotopic or non-isotopic labelling (by immunofluorescence or biotinilised probes), genetic amplification, (especially by PCR or LCR), double immunodiffusion technique, counter-electrophoresis technique, haemagglutination or a mixture thereof.

Another aspect of the present invention concerns a diagnosis method wherein a biological sample from the patient, such as cephalo-rachidian fluid, serum, blood, plasma, urine, broncho-alveolar lavage, stomach lavage, etc., is isolated from the patient, and is put in contact with the diagnostic device according to the invention for the diagnosis or the monitoring of an injury or a disease, preferably a lung injury or an oxidative stress-related disorder, affected by the presence of pro-oxidant agent or oxidative stress such as specific cardiovascular diseases like arteriosclerosis, neurodegenerative disorders (Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis), apoptosis, inflammatory reactions, allergic reactions such as asthma, hay fever and eczema, high bone mass syndrome, osteopetrosis, osteoporosis-pseudoglioma syndrome, and Bardet-Biedl syndrome 1. Said diagnosis and monitoring upon one or more biological samples obtained from several tissues from the patient can be advantageously obtained by one or more of the methods above-described, which could be adapted



according to the specific biological sample by the person skilled in the art.

Therefore, the product according to the invention could be used as a marker for the above-
5 identified injuries, diseases or disorders in a broad spectrum of tissues as shown in the enclosed Figure 1.

A further aspect of the present invention is related to a pharmaceutical composition comprising a pharmaceutically acceptable carrier and an element selected
10 from the group consisting of the nucleotide sequence, the amino acid sequence of the polypeptide according to the invention, the inhibitor directed against said sequences and/or one or more portions thereof.

A last aspect of the present invention is
15 related to the use of the pharmaceutical composition according to the invention for the manufacture of a medicament in the treatment and/or the prevention of lung injuries and/or diseases or of oxidative stress-related disorders.

20 The present invention is also related to a prevention and/or treatment method of a patient, especially a human patient, preferably affected by lung injuries and/or diseases or by oxidative stress-related disorders, wherein a sufficient amount of the pharmaceutical
25 composition according to the invention is administered to said patient in order to treat, avoid and/or reduce the symptoms of said injuries and/or diseases.

Other injuries and/or diseases which can be prevented and/or treated are injuries and/or diseases
30 affected by the presence of pro-oxidant agents or oxidative stress, such as specific cardio-vascular diseases like arteriosclerosis, neurodegenerative disorders such as

Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, apoptosis and inflammatory reactions and some allergic reactions such as asthma, hay fever and eczema, high bone mass syndrome, osteopetrosis, 5 osteoporosis-pseudoglioma syndrome, and Bardet-Biedl syndrome 1.

The pharmaceutically acceptable carrier according to the invention is any compatible non-toxic substance suitable for administering the composition 10 according to the invention to a human patient. Pharmaceutically acceptable carriers according to the invention suitable for oral administration are the ones well known by the person skilled in the art, such as tablets, coated or non-coated pills, capsules, spray-gas, 15 patches, gels, solutions or syrups. Pharmaceutically acceptable carriers vary according to the mode of administration (intravenous, intramuscular, subcutaneous, parenteral, etc.), and may comprise also adjuvants well known by the person skilled in the art to increase, reduce 20 and/or regulate humoral, local and/or cellular response of the immune system.


The pharmaceutical composition according to the invention may be prepared by the methods, generally applied by the person skilled in the art in the preparation 25 of various pharmaceutical compositions, wherein the percentage of the active compound/pharmaceutically acceptable carrier can vary within very large ranges, only limited by the tolerance of the patient to said pharmaceutical composition, and wherein the limits are 30 particularly determined by the frequency of administration and the possible side-effects of the active compounds or its pharmaceutically acceptable carrier.

Another aspect of the invention is related to the use of the diagnostic device according to the invention for performing upon the patient or upon a biological fluid obtained from the patient, a diagnosis, a dosage and/or a
5 monitoring of the above-mentioned injuries or diseases or oxidative stress-related disorders affecting the patient.

A further aspect of the present invention is related to a cell or a non-human animal, preferably a mammal such as a mouse or a rat, transformed by the vector
10 according to the invention and overexpressing the polypeptide according to the invention, or a non-human animal, preferably a mammal such as a mouse or a rat, genetically modified by a partial or total deletion of its genomic sequence encoding the polypeptide according to the
15 invention (knock-out non-human mammal) and obtained by methods well known by the person skilled in the art such as the one described by Kahn et al. (*Cell*, Vol. 92, pp. 593-596 (March 1998)).

Other examples of genetically modified non-
20 human animals according to the invention may be a transgenic non-human animal comprising an inhibitor according to the invention, preferably an antisense nucleic acid sequence complementary to the nucleotide sequence according to the invention so placed as to be transcribed
25 into antisense mRNA which is complementary to the nucleotide sequence according to the invention and which hybridises to said nucleotide sequence, thereby reducing or blocking its translation.

Further aspects of the present invention will
30 be described in the enclosed non-limiting examples in reference to the following Figures.



Brief description of the drawings

- Figure 1 represents a dot blot analysis of mRNA encoding the polypeptide according to the invention in various types of human tissues.
- 5 Figure 2 represents a Northern blot analysis of mRNA encoding the polypeptide according to the invention in a rat lung after administration of lipopolysaccharides (LPS) inducing an inflammatory reaction of the lung.
- 10 Figure 3 represents a Northern blot analysis of mRNA encoding the polypeptide according to the invention in a rat lung after intraperitoneal injection of pneumotoxigants.
- Figure 4 is a schematic representation of the human genomic sequence, the complete cDNA sequence and the corresponding amino acid sequence.
- 15 Figure 5 represents respectively the alignment of the sequences of the human B18 polypeptide according to the invention with the corresponding rat and mouse sequences.
- 20

Example 1 : Homology between the B18 polypeptide according to the invention with other known nucleotide or amino acid sequences

- 25 The BLAST 2.0 software (gapped BLAST at the NCBI Internet site) was used for searching for homologies between human B18 (162 amino acids) and known polypeptides in databases (GenBank, SwissProt). Said search did not give perfect alignment with known peptides from different
- 30 species (Table 1). Homologies of the human B18 cDNA (805 nucleotides) with GenBank, EMBL, DDBJ and PDB deposited

nucleotide sequences (Table 2) and GenBank Expression Sequence TAGS (ESTs) were noted.

Table 1 : Homologies of the B18 proteins (162 amino acid) with other proteins

| Name | NCBI ID | Identity (%) Homology (%) |
|--|----------|------------------------------|
| Membrane protein (synechocystis sp.) | 1652859 | 57/129 (44%) 81/129 (62%) |
| Peroxisomal-like protein (Aspergillus fumigatus) | 2769700 | 56/176 (31%) 90/176 (50%) |
| Haein HI0572 hypothetical protein (Haemophilus influenzae) | 1723174 | 53/146 (36%) 80/146 (54%) |
| PMP20 (Schizosaccharomyces pombe) | AJ002536 | 54/161 (33%) 85/161 (52%) |
| Peroxisomal membrane protein A (PMP 20) (Candida boidinii) | 130360 | 59/170 (34%) 89/170 (51%) |
| Peroxisomal membrane protein B (PMP 20) (Candida boidinii) | 130361 | 58/170 (34%) 88/170 (51%) |
| Putative peroxisomal protein PMP from yeast (Saccharomyces cerevisiae) | 1709682 | 41/138 (29%) 72/138 (51%) |
| Alkylhydroperoxide reductase C22 protein (Escherichia coli) | P26427 | 36/126 (28%) 58/126 (45%) |

Table 2

| Name | Access NO | Identity |
|--|-----------|---------------|
| Human mRNA down-regulated in cells infected by adenovirus 5 | U82616 | 259/263 (98%) |
| Human mRNA down-regulated in cells infected by adenovirus 5 | U82615 | 300/321 (93%) |

In the Table 2, an identity of 98% has been obtained with the alignment of 259 nucleotides of CDNA B18, which comprises in its totality 805 nucleotides, with 263 nucleotides of U82616 CDNA. A similar identity has been
5 obtained with the U82615 sequence.

The sequence SEQ ID NO 1 comprising 805 nucleotides presents a homology with several EST sequences obtained from a human and from a mouse, having the following references :

10 Human :

AA130751, N42215, W38597, N91311, N68467, AA187737,
N68916, W00593, R88950, AA181884, H20154, H66666

Mouse :

AA220019, AA123351, AA087129, AA255021, AA249897, W71344

15

Example 2 : Tissue detection

A human RNA master Blot (Clontech) containing 100-500 ng of poly-A + human RNA in each dot (normalised to the mRNA expression levels of eight different housekeeping
20 genes) was hybridised with a 554 bp-long B18 probe labelled with ³²P, and quantified using Phosphorimaging Technology. As shown in Figure 1, B18 mRNA is present in all tissues examined but predominantly in trachea, lung, kidney, thyroid gland, stomach, colon, heart and some regions of
25 the brain. Highest expression has been noted in the thyroid tissue. This presence is probably correlated with the possible antioxidant activity of the B18 polypeptide according to the invention.

30 Example 3 : Inflammatory reaction

Figure 2 represents a Northern blot analysis of rat lung mRNA after 6, 48 and 72 hours after

lipopolysaccharides (LPS) instillation inducing an inflammatory reaction in the lung.

A Northern blot containing 15 μ g of total RNA in each lane was hybridised with a 225 bp-long rat B18 probe, stripped and reprobed with a 572 bp-long rat β -actin probe, both labelled with 32 P. Northern blot was quantified using Phosphorimaging Technology and the B18 mRNA data were normalised to β -actin mRNA level.

10 Example 4 : Pneumotoxic reaction

Figure 3 represents a Northern blot analysis of rat lung mRNA after intraperitoneal injection of pneumotoxicants (4-ipomeanol, 1-(3-furyl)-4-hydroxypentanone (IPO), methylcyclopentadienyl manganese tricarbonyl (MMT) or alpha naphthylthiourea (ANTU)). These agents are known to induce in the lung acute lesions of Clara (IPO) and alveolar cells (MMT) as well as increasing the permeability of the alveolar/blood barrier (ANTU). A Northern blot containing 15 μ g of total RNA in each lane was hybridised with a 225 bp-long rat B18 probe, stripped and reprobed with a 572 bp-long β -actin probe both labelled with 32 P. The Northern blot was quantified using Phosphorimaging Technology and rat B18 mRNA data were normalised to β -actin mRNA level.

25

Example 5 : Proteinic activity of the B18 polypeptide

An amino analysis of the complete human B18 amino acid sequence shows that said polypeptide presents specific portions showing an homology with other anti-oxidant enzymes (starting from a Leucine at position 36 until a Cysteine at position 47) and an other portion

having an important homology with beta chains of ATP synthase (starting from a Glutamic acid at position 13 until a Glycine in position 38).


Furthermore, the B18 amino acid sequence according to the invention shows an important homology with an *Aspergillus fumigatus* allergen (34% identity and 60% homology by using clustal V sequence alignment), especially upon the portion of said B18 polypeptide having possible antioxidant properties. Therefore, it is possible that a peroxisomal protein (possibly homologous to B18 protein) is able to induce and to bind IgE from patients sensitised to *Aspergillus fumigatus* peroxisomal proteins after an induction of the patient immune system with *Aspergillus fumigatus* allergen. This mechanism can be compared to a reaction obtained with the manganese superoxide dismutase (MnSOD) wherein the human MnSOD is able to bind to IgE from patients sensitised to *Aspergillus fumigatus* MnSOD.

Furthermore, the Inventors have identified a portion of the B18 human polypeptide which presents an homology with a Cyclophilin-binding domain of *Candida boidinii* PMP20 (receptor, of the immuno-suppressant drug cyclosporine A). Said possible Cyclophilin-binding domain is starting from the Threonine in position 150 until the Leucine in position 161.

25

Example 6 : B18 human gene and mRNA alternative splicing

As represented in the enclosed Figure 4, the Inventors have identified upon the genomic DNA (SEQ ID NO 10) 5 exons and 5 introns. By RT-PCR (using primers 5'-gggtatgggactagctggcg-3' and 5'-ctggccaacattccaattgcag-3') and according to the genomic sequence, 4 different cDNAs corresponding to the transcription of the said genomic DNA



have been identified in human lung and in human brain. A first cDNA of 736 bp corresponds to the cDNA encoding the complete amino acid sequence of the B18 protein according to the invention. However, 3 other cDNAs of 601, 604 and 5 469 bp were also identified, and comprise specific splicings of one or more exons.

Therefore, another aspect of the present invention is related to said specific portions of the complete genomic or CDNA nucleotide sequence according to 10 the invention or to specific portions of the complete amino acid sequence of the B18 protein according to the invention, which could be used also as specific markers of the B18 activity, preferably the anti-oxidative activity.

15 **Example 7 : Knock-out mouse**

Exons of a mouse genomic sequence encoding the B18 polypeptide according to the invention have been deleted by homologous recombination. Said homologous recombination has been obtained with a genetic sequence 20 comprising a neomycin resistant gene. The targeting vector with said gene and a ,thymidine kinase (in order to eliminate non-homologous recombinants with ganciclovir) has been prepared. Said recombination was used for the deletion of one or more exons of the B18 polypeptide. After 25 electroporation of ES cells with the targeting vector, positive clones having incorporated homologous recombination were identified by Southern blot with labelled probes. Aggregation of said positive clones with a morula from a Swiss pseudo-pregnant mouse produces several 30 chimeric mice which survive after birth. Several homozygote mice are obtained by cross-breeding and are used as a model for the above-mentioned diseases.

Similar experiments may be done with another mammal whose B18 sequence is known (the B18 sequence of a mouse and a rat and their alignment with the human sequence is shown in the enclosed Figure 5).

5


Example 8 : Chromosome localisation of human B18 gene

Radiation hybrid clones (GeneBridge 4 Radiation Hybrid Panel, Research Genetics) were used for performing chromosome localisation by PCR with two
10 different pairs of primers (5'-caggttcaccttggtccctggctc-3' (SEQ ID NO 14), 5'-atgttatgcaaccctttgcgacac-3' (SEQ ID NO 17) and 5'-gtgtttgaaggggagccaggggaac-3' (SEQ ID NO 18), 5'-agagacagggtttcaccatcttg-3' (SEQ ID NO 19)).

The Inventors have located B18 genomic
15 sequence on human chromosome 11q13. B18 gene has been located 7.15-6.1 cR from marker D11S913 between markers D11S1963 and D11S4407 (Genome Database internet site).

Unknown genes linked to different disorders have been localised in the same region of chromosome 11.
20 Therefore, B18 gene is possibly associated with these disorders:

- atopy (atopic hypersensitivity: asthma, hay fever and eczema; MIM n°147050 at OMIM of NCBI internet site),
- high bone mass syndrome (MIM n°601884),
- 25 - osteopetrosis (MIM n°259700),
- osteoporosis-pseudoglioma syndrome (MIM n°259770) and
- Bardet-Biedl syndrome 1 (MIM n°209901).



CLAIMS

1. Amino acid sequence having more than 70% homology with the sequence SEQ ID NO 2.
2. Amino acid sequence according to claim 1,
5 having more than 85% homology with the sequence SEQ ID NO 2.
3. Amino acid sequence according to claim 1 or 2, having more than 95% homology with the sequence SEQ ID NO 2.
- 10 4. Amino acid sequence according to any one of the preceding claims, corresponding to SEQ ID NO 2 or an immunoreactive portion thereof.
5. Nucleotide sequence encoding the amino acid sequence according to any one of the preceding claims
15 and presenting more than 70% homology with SEQ ID NO 1 or its complementary strand.
6. Nucleotide sequence according to claim 5, having more than 85% homology with the sequence SEQ ID NO 1 or its complementary strand.
- 20 7. Nucleotide sequence according to claim 5 more than 95% homology with the sequence SEQ ID NO 1 or its complementary strand.
8. Nucleotide sequence according to any one of the claims 5 to 7, corresponding to the sequence SEQ ID
25 NO 1, its complementary strand or a portion thereof specific for SEQ ID NO 1 and comprising more than 15 base pairs.
9. Vector comprising the nucleotide sequence according to any one of the claims 5 to 8.
- 30 10. Inhibitor directed against the amino acid or nucleotide sequence according to any one of the claims 1 to 8.

11. Inhibitor according to claim 10, being an antibody, preferably a monoclonal antibody, or a portion of said antibody.

12. Diagnostic device comprising an element
5 selected from the group consisting of the amino acid sequence according to any one of the claims 1 to 4, the nucleotide sequence according to any one of the claims 5 to 8, the inhibitor according to claim 10 or 11, their portions or a mixture thereof.

10 13. Method for the in vitro detection of lung injuries and diseases or oxidative stress-related diseases and disorders, especially inflammatory diseases, comprising the steps of :

- isolating a sample from a body fluid of a patient,
15 preferably a human patient,
- possibly inhibiting the contaminants present in said sample,
- put in contact said sample with an element selected from the group consisting of the amino acid sequence
20 according to any one of the claims 1 to 4, the nucleotide sequence according to any one of the claims 5 to 8, the inhibitor according to claim 10 or 11, their portions or a mixture thereof, and
- detecting a reaction of a molecule present in said
25 sample with said element.

14. Pharmaceutical composition comprising a pharmaceutically acceptable carrier and an element selected from the group consisting of the amino acid sequence according to any one of the claims 1 to 4, the nucleotide
30 sequence according to any one of the claims 5 to 8, the

inhibitor according to claim 10 or 11, their portions or a mixture thereof.

15. Use of the pharmaceutical composition according to claim 14 for the manufacture of a medicament
5 for the prevention and/or the treatment of lung injuries or diseases, and of oxidative stress-related diseases or disorders, such as specific cardio-vascular diseases like arteriosclerosis, neurodegenerative disorders such as Alzheimer's disease, Parkinson's disease, amyotrophic
10 lateral sclerosis, apoptosis and inflammatory reactions, allergic reactions such as asthma, hay fever and eczema, high bone mass syndrome, osteopetrosis, osteoporosis-pseudoglioma syndrome, and Bardet-Biedl syndrome 1.

16. Cell transformed by the vector according
15 to claim 9 or comprising a partial or total deletion of its nucleotide sequence according to any one of the claims 5 to 8.

17. Non-human animal, preferably a mammal,
transformed by the vector according to claim 9 or
20 comprising a partial or total deletion of its nucleotide sequence according to any, one of the claims 5 to 8.

1/7







FIG.1




2/7




3/7



Gene (chromosome 11q12-13)



mRNAs



Proteins




FIG. 4

CLUSTAL V alignment of human and rat B18 amino acid sequences (Identity: 90%, Homology: 97.5%):

```

B18hum      MAPIKVGDAIPAVEVFEGEPGNKVNLAELFKGKKGVLFVPGAFTPGCSK = SEQIDNO1
B18rat      MAPIKVGDTIPSVEVFEGEPGKKVNLAELFKDKKGVLFVPGAFTPGCSK
*****
B18hum      THLPGFVEQAEALKAKGVQVVACLSVNDAFVTGEWGRAHKAEGKVRLAD
B18rat      THLPGFVEQAGALKAKGAQVVACLSVNDVFVTAEWGRAHQAEGKVQLLAD
*****
B18hum      PTGAFGKETDLLLDDSLVSI FGNRRLLKRFSMVVDGIVKALNVEPDGTGL
B18rat      PTGAFGKETDLLLDDSLVSLFGNRRLLKRFSMVIDKGVVKALNVEPDGTGL
*****
B18hum      TCSLAPNIISQL
B18rat      TCSLAPNILSQL
*****

```

FIG.5a

CLUSTAL V alignment of human and mouse B18 amino acid sequences (Identity: 91%, Homology: 96%):

```

B18hum      MAPIKVGDAIPAVEVFEGEPGNKVNLAELFKGKKGVLFVPGAFTPGCSK
B18mouse    MAPIKVGDAIPSVEVFEGEPGKKVNLAELFKGKKGVLFVPGAFTPGCSK
*****
B18hum      THLPGFVEQAEALKAKGVQVVACLSVNDAFVTGEWGRAHKAEGKVRLAD
B18mouse    THLPGFVEQAGALKAKGAQVVACLSVNDVFIEEWGRAHQAEGKVRLAD
*****
B18hum      PTGAFGKETDLLLDDSLVSI FGNRRLLKRFSMVVDGIVKALNVEPDGTGL
B18mouse    PTGAFGKATDLLLDDSLVSLFGNRRLLKRFSMVIDNGIVKALNVEPDGTGL
*****
B18hum      TCSLAPNIISQL
B18mouse    TCSLAPNILSQL
*****

```

CLUSTAL V alignment of human and rat cDNA sequences (identity: 612/780, 78.5%):

```

B18hum      GCCAGGAGGCGGAGTGGAGTGGCCGTGGGGCGGGTATGGGACTAGCTGG
B18rat      -----TG-----CGTC-----CTAGGCAG
                **          ***          ****
B18hum      CGTGTGCGCCCTGAGACGCTCAGCGGGCTATATACTCGTCGGTGGGGCCG
B18rat      CATA---GCC---GGA---TCGGTGCTCCGTGCATCGGCTACTTGGAC--
                * *      ***      * * * * * * * * * *
B18hum      GCGGTCAGTCTGCGGCAGCGGCAGCAAGACGGTGCAGTGAAGGAGAGTGG
B18rat      -----GTGCGTGGCAGGCAGCAGGCCGG---AAAGGAGCAGGTTGG
                * * * * * * * * * * * * * * * *

```


6/7

FIG. 5b

| | |
|--------|--|
| B18hum | CGCTCTGGCGGGGTCCGCAGTTTCAGCAGAGCCGCTGCAGCCATGGCCCC |
| B18rat | GAGTGTGGTGGGGCCCGCAGCTTCAGCAGTGGCGGGTGACTATGGCCCC * * * * * |
| B18hum | AATCAAGGTGGGAGATGCCATCCCAGCAGTGGAGGTGTTTGAAGGGGAGC |
| B18rat | GATCAAGGTGGGAGACACCATTCCCTCAGTGGAGGTATTTGAAGGGGAAC * * * * * |
| B18hum | CAGGGAACAAGGTGAACCTGGCAGAGCTGTTCAAGGGCAAGAAGGGTGTG |
| B18rat | CTGGAAAGAAGGTGAACCTGGCAGAGCTGTTCAAGGACAAGAAAGGTGTT * * * * * |
| B18hum | CTGTTTGGAGTTCCTGGGGCCTTCACCCCTGGATGTTCCAAGACACACCT |
| B18rat | TTGTTTGGAGTCCCTGGGGCATTACACCTGGCTGTTCCAAGACCCATCT * * * * * |
| B18hum | GCCAGGGTTTGTGGAGCAGGCTGAGGCTCTGAAGGCCAAGGGAGTCCAGG |
| B18rat | GCCTGGGTTTGTGGAGCAAGCCGGAGCTCTGAAGGCCAAGGGAGCACAAAG * * * * * |
| B18hum | TGGTGGCCTGTCTGAGTGTAAATGATGCCTTTGTGACTGGCGAGTGGGGC |
| B18rat | TGGTGGCCTGTCTGAGTGTAAATGATGTCTTCGTGACTGCAGAGTGGGGT * * * * * |
| B18hum | CGAGCCCACAAGGCGGAAGGCAAGGTTCTGGCTCCTGGCTGATCCCACTGG |
| B18rat | CGAGCCCACCAAGGCAAGGCAAGGTTCTGGCTCCTGGCTGACCCCACTGG * * * * * |
| B18hum | GGCCTTTGGGAAGGAGACAGACTTATTACTAGATGATTCGCTGGTGTCCA |
| B18rat | AGCTTTTGGAAAGGAGACAGATTTACTACTAGATGATTCCTTGGTGTCTC * * * * * |
| B18hum | TCTTTGGGAATCGACGTCTCAAGAGGTTCTCCATGGTGGTACAGGATGGC |
| B18rat | TCTTTGGGAATCGTCGGCTAAAAGGTTCTCCATGGTGATAGACAAGGGC * * * * * |
| B18hum | ATAGTGAAGGCCCTGAATGTGGAACCAGATGGCACAGGCCTCACCTGCAG |
| B18rat | GTAGTAAAGGCACTGAACGTGGAGCCGGATGGCACAGGCCTCACCTGCAG * * * * * |
| B18hum | CCTGGCACCCAATATCATCTCACAGCTCTGAGGCCCTGGGCCAGATTACT |
| B18rat | CCTGGCCCCCAACATCTCTCAAACTCTGAGGCCCTGA-CCAGA--ATG * * * * * |
| B18hum | TCCTCCACCCCTCCCTATCTCACCTGCCAGCCCTGTGCTGG-GGCCCTG |
| B18rat | TCCTCTGACTCTCC-ATCTCCTCCACCCAGCTCTGGGCCAAAGGCCAG * * * * * |
| B18hum | CA-----ATTGGAATG-----TTGGCCAGATTTCTGC |
| B18rat | TACCTCCTTACCTGAGGGCCACTGGAATGGAACCTTGACAATATTTCTGC * * * * * |
| B18hum | AATAAACACTTGTGGTTTGCAGAAAAA----- |
| B18rat | AATAAACAGTT-TAATTTGTGAAAAAAAAAAAAAAAAAAAA * * * * * |

CLUSTAL V alignment of human and mouse cDNA sequences (Identity: 552/675, 81.8%):

FIG. 5c

```

B18hum      GCCAGGAGGCGGAGTGGAAAGTGGCCGTGGGGCGGGTATGGGACTAGCTGG
B18mouse    -----

B18hum      CGTGTGCGCCCTGAGACGCTCAGCGGGCTATATACTCGTCGGTGGGGCCG
B18mouse    -----TGCTCCGTG-----CATCGACGTGCTTG
                  **** * * * * *

B18hum      GCGGTCTAGTCTGCGGCAGCGGCAGCAAGACGGTGCAGTGAAGGAGAGTGG
B18mouse    GCAGGCAG-----AGCAGGCCGG---AAAGAAGCAGGTTGG
                  * * * * *

B18hum      GCGTCTGGCGGGGTCCGCAGTTTCAGCAGAGCCGCTGCAGCCATGGCCCC
B18mouse    GAGTGTGGCGGAGCCCGCAGCTTCAGCAGCTCCGCGGTGACCATGGCCCC
                  * * * * *

B18hum      AATCAAGGTGGGAGATGCCATCCCAGCAGTGGAGGTGTTTGAAGGGGAGC
B18mouse    GATCAAGGTGGGAGATGCCATTCCCTCAGTGGAGGTATTTGAAGGGGAAC
                  * * * * *

B18hum      CAGGGAACAAGGTGAACCTGGCAGAGCTGTTCAAGGGCAAGAAGGGTGTG
B18mouse    CGGGAAGAAGGTGAACCTGGCAGAGCTGTTCAAGGGCAAGAAAGGTGTT
                  * * * * *

B18hum      CTGTTTGGAGTTCCTGGGGCCTTCACCCCTGGATGTTCCAAGACACACCT
B18mouse    TTGTTTGGAGTCCCTGGGGCATTACACCTGGCTGTTCTAAGACCCACCT
                  * * * * *

B18hum      GCCAGGGTTTGTGGAGCAGGCTGAGGCTCTGAAGGCCAAGGGAGTCCAGG
B18mouse    GCCTGGGTTTGTGGAGCAAGCTGGAGCTCTGAAGGCTAAGGGAGCGCAGG
                  * * * * *

B18hum      TGGTGGCCTGTCTGAGTGTTAATGATGCCTTTGTGACTGGCGAGTGGGGC
B18mouse    TGGTGGCCTGTCTGAGCGTTAATGACGTCTTTGTGATTGAAGAGTGGGGT
                  * * * * *

B18hum      CGAGCCCACAAGGCGGAAGGCAAGGTTGCGCTCCTGGCTGATCCCACTGG
B18mouse    CGAGCCCACCAGGCAGAAGGCAAGGTTGCGCTCCTGGCTGACCCCACTGG
                  * * * * *

B18hum      GGCCTTTGGGAAGGAGACAGACTTATTACTAGATGATTGCTGGTGTCCA
B18mouse    AGCCTTTGGGAAGGCGACAGACTTATTATTGGATGATTCTTTGGTGTCTC
                  * * * * *

B18hum      TCTTTGGGAATCGACGTCTCAAGAGGTTCTCCATGGTGGTACAGGATGGC
B18mouse    TCTTTGGGAATCGTCGGCTGAAAAGGTTCTCCATGGTGGTACAGACAACGGC
                  * * * * *

B18hum      ATAGTGAAGGCCCTGAATGTGGAACCAGATGGCACAGGCCTCACCTGCAG
B18mouse    ATAGTGAAGGCACTGAACGTGGAGCCAGATGGCACAGGCCTCACCTGCAG
                  * * * * *

B18hum      CCTGGCACCCAATATCATCTCACAGCTCTGAGGCCCTGGGCCAGATTACT
B18mouse    CCTGGCCCCCAACATCCTCTCCCAACTCTGAGGCCCTGG-CCAGATG---
                  * * * * *

B18hum      TCCTCCACCCCTCCCTATCTCACCTGCCCAGCCCTGTGCTGGGGCCCTGC
B18mouse    TCCTCTGACTCTCC-ATCTCTCCACCCGGCTCT-----AGGCC---
                  * * * * *

B18hum      AATTGGAATGTTGGCCAGATTCTGCAATAAACACTTGTGGTTTGGCGAA
B18mouse    ----AAAAGGCTCGGTACCTCCTTACTGGGAGC-CACGT-----
                  * * * * *

```


1
SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: UNIVERSITE CATHOLIQUE DE LOUVAIN
Halles Universitaires
(B) STREET: Place de l' Universite, 1
(C) CITY: LOUVAIN-LA-NEUVE
(E) COUNTRY: BELGIUM
(F) POSTAL CODE (ZIP): B-1348

(A) NAME: UNIVERSITE DE MONS-HAINAUT
(B) STREET: Place du Parc 20
(C) CITY: MONS
(E) COUNTRY: BELGIUM
(F) POSTAL CODE (ZIP): B-7000

(ii) TITLE OF INVENTION: PEROXISOME-ASSOCIATED PEPTIDE, NUCLEOTIDE
SEQUENCE ENCODING SAID PEPTIDE AND THEIR USES IN THE
DIAGNOSTIC AND/OR THE TREATMENT OF LUNG INJURIES AND
DISEASES, AND OF OXIDATIVE STRESS-RELATED DISORDERS

(iii) NUMBER OF SEQUENCES: 19

(iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 805 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 193..681

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

```
GCCAGGAGGC GGAGTGGAAG TGGCCGTGGG GCGGGTATGG GACTAGCTGG CGTGTGCGCC      60
CTGAGACGCT CAGCGGGCTA TATACTCGTC GGTGGGGCCG GCGGTCAGTC TCGGGCAGCG      120
```


2

| | |
|--|-----|
| GCAGCAAGAC GGTGCACTGA AGGAGAGTGG GCGTCTGGCG GGGTCCGCAG TTTCAGCAGA | 180 |
| GCCGCTGCAG CC ATG GCC CCA ATC AAG GTG GGA GAT GCC ATC CCA GCA Met Ala Pro Ile Lys Val Gly Asp Ala Ile Pro Ala 1 5 10 | 228 |
| GTG GAG GTG TTT GAA GGG GAG CCA GGG AAC AAG GTG AAC CTG GCA GAG Val Glu Val Phe Glu Gly Glu Pro Gly Asn Lys Val Asn Leu Ala Glu 15 20 25 | 276 |
| CTG TTC AAG GGC AAG AAG GGT GTG CTG TTT GGA GTT CCT GGG GCC TTC Leu Phe Lys Gly Lys Lys Gly Val Leu Phe Gly Val Pro Gly Ala Phe 30 35 40 | 324 |
| ACC CCT GGA TGT TCC AAG ACA CAC CTG CCA GGG TTT GTG GAG CAG GCT Thr Pro Gly Cys Ser Lys Thr His Leu Pro Gly Phe Val Glu Gln Ala 45 50 55 60 | 372 |
| GAG GCT CTG AAG GCC AAG GGA GTC CAG GTG GTG GCC TGT CTG AGT GTT Glu Ala Leu Lys Ala Lys Gly Val Gln Val Val Ala Cys Leu Ser Val 65 70 75 | 420 |
| AAT GAT GCC TTT GTG ACT GGC GAG TGG GGC CGA GCC CAC AAG GCG GAA Asn Asp Ala Phe Val Thr Gly Glu Trp Gly Arg Ala His Lys Ala Glu 80 85 90 | 468 |
| GGC AAG GTT CGG CTC CTG GCT GAT CCC ACT GGG GCC TTT GGG AAG GAG Gly Lys Val Arg Leu Leu Ala Asp Pro Thr Gly Ala Phe Gly Lys Glu 95 100 105 | 516 |
| ACA GAC TTA TTA CTA GAT GAT TCG CTG GTG TCC ATC TTT GGG AAT CGA Thr Asp Leu Leu Leu Asp Asp Ser Leu Val Ser Ile Phe Gly Asn Arg 110 115 120 | 564 |
| CGT CTC AAG AGG TTC TCC ATG GTG GTA CAG GAT GGC ATA GTG AAG GCC Arg Leu Lys Arg Phe Ser Met Val Val Gln Asp Gly Ile Val Lys Ala 125 130 135 140 | 612 |
| CTG AAT GTG GAA CCA GAT GGC ACA GGC CTC ACC TGC AGC CTG GCA CCC Leu Asn Val Glu Pro Asp Gly Thr Gly Leu Thr Cys Ser Leu Ala Pro 145 150 155 | 660 |
| AAT ATC ATC TCA CAG CTC TGA GGCCCTGGGC CAGATTACTT CCTCCACCCC Asn Ile Ile Ser Gln Leu * 160 | 711 |
| TCCCTATCTC ACCTGCCCAG CCCTGTGCTG GGGCCCTGCA ATTGGAATGT TGGCCAGATT | 771 |
| TCTGCAATAA ACACTTGTGG TTTGCGGAAA AAAA | 805 |

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 163 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

3

Met Ala Pro Ile Lys Val Gly Asp Ala Ile Pro Ala Val Glu Val Phe
 1 5 10 15

Glu Gly Glu Pro Gly Asn Lys Val Asn Leu Ala Glu Leu Phe Lys Gly
 20 25 30

Lys Lys Gly Val Leu Phe Gly Val Pro Gly Ala Phe Thr Pro Gly Cys
 35 40 45

Ser Lys Thr His Leu Pro Gly Phe Val Glu Gln Ala Glu Ala Leu Lys
 50 55 60

Ala Lys Gly Val Gln Val Val Ala Cys Leu Ser Val Asn Asp Ala Phe
 65 70 75 80

Val Thr Gly Glu Trp Gly Arg Ala His Lys Ala Glu Gly Lys Val Arg
 85 90 95

Leu Leu Ala Asp Pro Thr Gly Ala Phe Gly Lys Glu Thr Asp Leu Leu
 100 105 110

Leu Asp Asp Ser Leu Val Ser Ile Phe Gly Asn Arg Arg Leu Lys Arg
 115 120 125

Phe Ser Met Val Val Gln Asp Gly Ile Val Lys Ala Leu Asn Val Glu
 130 135 140

Pro Asp Gly Thr Gly Leu Thr Cys Ser Leu Ala Pro Asn Ile Ile Ser
 145 150 155 160

Gln Leu *

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 780 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Rattus Rattus

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 136..624

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

TGCGTCCTAG GCAGCATAGC CGGATCGGTG CTCCGTGCAT CGGCTACTTG GACGTGCGTG 60

GCAGGCAGAG CAGGCCGGAA AGGAGCAGGT TGGGAGTGTG GTGGGGCCCG CAGCTTCAGC 120

4

```

AGTGCCGCGG TGA CTATGGC CCCGATCAAG GTGGGAGACA CCATTCCCTC AGTGGAGGTA      180
TTTGRAGGGG AACCTGGAAA GAAGGTGAAC TTGGCAGAGC TGTTC AAGGA CAAGAAAGGT      240
GTTTTGTTTG GAGTCCCTGG GGCATTTACA CCTGGCTGTT CCAAGACCCA TCTGCCTGGG      300
TTTGTGGAGC AAGCCGGAGC TCYGAAGGCC AAGGGAGCAC AAGTGGTGGC CTGTCTGAGT      360
GTTAATGATG YCTTCGTGAC TGCAGAGTGG GGTGAGCCC ACCAGGCAGA AGGCAAGGTT      420
CAGCTCCTGG CTGACCCAC TGGAGCTTTT GGAAAGGAGA CAGATTTACT ACTAGATGAT      480
TCTTTGGTGT CTCTCTTTGG GAATCGTCGG CTAAAAAGGT TCTCCATGGT GATAGACAAG      540
GGCGTAGTAA AGGCACTGAA CGTGGAGCCG GATGGCACAG GCCTCACCTG CAGCCTGGCC      600
CCCAACATCC TCTCACA ACT CTGAGGCCCT GACCAGAATG TCCTCTGACT CTCCCATCTC      660
CTCCACCCAG CTCTGGGCCA AAGGCCAGT ACCTCCTTAC CTGAGGGCCA CTGGAATGGA      720
ACCTTGACAA TATTTCTGCA ATAAACAGTT TAATTTGTGA AAAAAAAAAA AAAAAAAAAA      780

```

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 162 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Rattus Rattus

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 17
- (D) OTHER INFORMATION: /product= "Glu or Gly"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 63
- (D) OTHER INFORMATION: /product= "Leu or Pro"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 79
- (D) OTHER INFORMATION: /product= "Ala or Val"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

```

Met Ala Pro Ile Lys Val Gly Asp Thr Ile Pro Ser Val Glu Val Phe
1           5           10           15

```


5

Xaa Gly Glu Pro Gly Lys Lys Val Asn Leu Ala Glu Leu Phe Lys Asp
 20 25 30

Lys Lys Gly Val Leu Phe Gly Val Pro Gly Ala Phe Thr Pro Gly Cys
 35 40 45

Ser Lys Thr His Leu Pro Gly Phe Val Glu Gln Ala Gly Ala Xaa Lys
 50 55 60

Ala Lys Gly Ala Gln Val Val Ala Cys Leu Ser Val Asn Asp Xaa Phe
 65 70 75 80

Val Thr Ala Glu Trp Gly Arg Ala His Gln Ala Glu Gly Lys Val Gln
 85 90 95

Leu Leu Ala Asp Pro Thr Gly Ala Phe Gly Lys Glu Thr Asp Leu Leu
 100 105 110

Leu Asp Asp Ser Leu Val Ser Leu Phe Gly Asn Arg Arg Leu Lys Arg
 115 120 125

Phe Ser Met Val Ile Asp Lys Gly Val Val Lys Ala Leu Asn Val Glu
 130 135 140

Pro Asp Gly Thr Gly Leu Thr Cys Ser Leu Ala Pro Asn Ile Leu Ser
 145 150 155 160

Gln Leu

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 675 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Mouse

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 99..588

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

| | |
|---|-----|
| TGCTCCGTGC ATCGACGTGC TTGGCAGGCA GAGCAGGCCG GAAAGAAGCA GGTGGGGAGT | 60 |
| GTGGCGGAGC CCGCAGCTTC AGCAGCTCCG CGGTGACCAT GGCCCCGATC AAGGTGGGAG | 120 |
| ATGCCATTCC CTCAGTGGAG GTATTTGAAG GGAACCGGG AAAGAAGGTG AACTTGGCAG | 180 |
| AGCTGTTCAA GGGCAAGAAA GGTGTTTTGT TTGGAGTCCC TGGGGCATT ACACCTGGCT | 240 |


```

GTTCTAAGAC CCACCTGCCT GGGTTTGTGG AGCAAGCTGG AGCTCTGAAG GCTAAGGGAG      300
CGCAGGTGGT GGCCTGTCTG AGCGTTAATG ACGTCTTTGT GATTGAAGAG TGGGGTCGAG      360
CCCACCAGGC AGAAGGCAAG GTTCGGCTCC TGGCTGACCC CACTGGAGCC TTTGGGAAGG      420
CGACAGACTT ATTATTGGAT GATTCTTTGG TGTCTCTCTT TGGGAATCGT CGGCTGAAAA      480
GGTTCTCCAT GGTGATAGAG AACGGCATAG TGAAGGCACT GAACGTGGAG CCAGATGGCA      540
CAGGCCTCAC CTGCAGCCTG GCCCCAACA TCCTCTCCA ACTCTGAGGC CCTGGCCAGA      600
TGTCTCTGA CTCTCCCATC TCTCCCACCC GGCTCTAGGC CAAAAGGCTC GGTACCTCCT      660
TACTGGGAGC CACGT                                                    675

```

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 162 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Mouse

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

```

Met Ala Pro Ile Lys Val Gly Asp Ala Ile Pro Ser Val Glu Val Phe
 1             5             10             15

Glu Gly Glu Pro Gly Lys Lys Val Asn Leu Ala Glu Leu Phe Lys Gly
      20             25             30

Lys Lys Gly Val Leu Phe Gly Val Pro Gly Ala Phe Thr Pro Gly Cys
      35             40             45

Ser Lys Thr His Leu Pro Gly Phe Val Glu Gln Ala Gly Ala Leu Lys
      50             55             60

Ala Lys Gly Ala Gln Val Val Ala Cys Leu Ser Val Asn Asp Val Phe
      65             70             75             80

Val Ile Glu Glu Trp Gly Arg Ala His Gln Ala Glu Gly Lys Val Arg
      85             90             95

Leu Leu Ala Asp Pro Thr Gly Ala Phe Gly Lys Ala Thr Asp Leu Leu
      100            105            110

Leu Asp Asp Ser Leu Val Ser Leu Phe Gly Asn Arg Arg Leu Lys Arg
      115            120            125

```


Phe Ser Met Val Ile Asp Asn Gly Ile Val Lys Ala Leu Asn Val Glu
 130 135 140

Pro Asp Gly Thr Gly Leu Thr Cys Ser Leu Ala Pro Asn Ile Leu Ser
 145 150 155 160

Gln Leu

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 469 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 161..382

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

```

GGGTATGGGA CTAGCTGGCG TGTGCGCCCT GAGACGCTCA GCGGGCTATA TACTCGTCGG      60
TGGGGCCGGC GGTCACTCTG CGGCAGCGGC AGCAAGACGG TGCAGTGAAG GAGAGTGGGC      120
GTCTGGCGGG GTCCGCAGTT TCAGCAGAGC CGCTGCAGCC ATGGCCCCAA TCAAGGTTCC      180
GCTCCTGGCT GATCCCACTG GGGCCTTTGG GAAGGAGACA GACTTATTAC TAGATGATTC      240
GCTGGTGTCC ATCTTTGGGA ATCGACGTCT CAAGAGGTTT TCCATGCTGG TACAGGATGG      300
CATAGTGAAG GCCCTGAATG TGAACACAGA TGGCACAGGC CTCACCTGCA GCCTGGCACC      360
CAATATCATC TCACAGCTCT GAGGCCCTGG GCCAGATTAC TTCCTCCACC CCTCCCTATC      420
TCACCTGCCC AGCCGTGTGC TGGGGCCCTG CAATTGGAAT GTTGCCAG      469
  
```

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 601 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION:161..514

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

```
GGGTATGGGA CTAGCTGGCG TGTGCGCCCT GAGACGCTCA GCGGGCTATA TACTCGTCGG      60
TGGGGCCGGC GGTCACTCTG CGGCAGCGGC AGCAAGACGG TGCAGTGAAG GAGAGTGGGC      120
GTCTGGCGGG GTCCGCAGTT TCAGCAGAGC CGCTGCAGCC ATGGCCCCAA TCAAGACACA      180
CCTGCCAGGG TTTGTGGAGC AGGCTGAGGC TCTGAAGGCC AAGGGAGTCC AGGTGGTGGC      240
CTGTCTGAGT GTTAATGATG CCTTTGTGAC TGGCGAGTGG GGCCGAGCCC ACAAGGCGGA      300
AGGCAAGGTT CGGCTCCTGG CTGATCCCAC TGGGGCCTTT GGAAGGAGA CAGACTTATT      360
ACTAGATGAT TCGCTGGTGT CCATCTTTGG GAATCGACGT CTCAAGAGGT TCTCCATGGT      420
GGTACAGGAT GGCATAGTGA AGGCCCTGAA TGTGGAACCA GATGGCACAG GCCTCACCTG      480
CAGCCTGGCA CCCAATATCA TCTCACAGCT CTGAGGCCCT GGGCCAGATT ACTTCCTCCA      540
CCCCCTCCCTA TCTCACCTGC CCAGCCCTGT GCTGGGGCCC TGCAATTGGA ATGTTGGCCA      600
G                                                                           601
```

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 604 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION:161..517

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

```
GGGTATGGGA CTAGCTGGCG TGTGCGCCCT GAGACGCTCA GCGGGCTATA TACTCGTCGG      60
```


9

| | |
|---|-----|
| TGGGGCCGGC GGTCACTCTG CGGCAGCGGC AGCAAGACGG TGCAGTGAAG GAGAGTGGGC | 120 |
| GTCTGGCGGG GTCCGCAGTT TCAGCAGAGC CGCTGCAGCC ATGGCCCCAA TCAAGGTGGG | 180 |
| AGATGCCATC CCAGCAGTGG AGGTGTTTGA AGGGGAGCCA GGAACAAGG TGAACCTGGC | 240 |
| AGAGCTGTTT AAGGGCAAGA AGGGTGTGCT GTTTGGAGTT CCTGGGGCCT TCACCCCTGG | 300 |
| ATGTTCCAAG GTTCGGCTCC TGGCTGATCC CACTGGGGCC TTTGGGAAGG AGACAGACTT | 360 |
| ATTACTAGAT GATTCGCTGG TGTCCATCTT TGGGAATCGA CGTCTCAAGA GGTCTCCAT | 420 |
| GGTGGTACAG GATGGCATAG TGAAGGCCCT GAATGTGGAA CCAGATGGCA CAGGCCTCAC | 480 |
| CTGCAGCCTG GCACCCAATA TCATCTCACA GCTCTGAGGC CCTGGGCCAG ATTACTTCCT | 540 |
| CCACCCCTCC CTATCTCACC TGCCAGCCC TGTGCTGGGG CCCTGCAATT GGAATGTTGG | 600 |
| CCAG | 604 |

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2710 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(ix) FEATURE:

- (A) NAME/KEY: exon
- (B) LOCATION:2516..2710

(ix) FEATURE:

- (A) NAME/KEY: exon
- (B) LOCATION:2074..2135

(ix) FEATURE:

- (A) NAME/KEY: exon
- (B) LOCATION:1932..1970

(ix) FEATURE:

- (A) NAME/KEY: exon
- (B) LOCATION:1728..1859

(ix) FEATURE:

- (A) NAME/KEY: exon
- (B) LOCATION:802..936

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

| | |
|--|------|
| TCTGTCCCTT AGCGCCCCCG CGGGGGCTTA CCCCATCCCA CTCCATGACC TCCCCTCCCC | 60 |
| CCATGGCGAA TTCCCACCTT TCTGTCTTTC ACTCACTTCC TGGAACCGTC CCCAGGGCCT | 120 |
| TGGACCTTCC CCCTTCTCCT CCCAAACCTT GTGAGACCCC ATTCCCTTTC TACTTCATCC | 180 |
| TGCTCTCAAC TTTTGGGCTC CTCAGAGGCC CTCACCCCTG ACTCTCTCTC CCTACCCACT | 240 |
| CTGGTCCCAT GAAGCCCTCA AGTACTCTGG GGATGGATCC TTCCCCCTTC AAAAGATTCC | 300 |
| TTCTTTTGT CTACACCTCC TGGGTGTAGG GGCCTGGACA CCCTCCCCCA ACGTTCCACC | 360 |
| TGCCGCTGCC CTTCTCTTTC CTCCTCCTGA GGGTGGGACC CTCAGACCTG GCCAAGATCC | 420 |
| TCTCCCTCCA TGTGTGTCAGG GACTCCTCCT CACCCCCAAA TACAGCCCTC TAGCCCCTGT | 480 |
| CCATTTTATT CCACTCCTTT CCTGTAACCT AGACAGCATG TTATGCAACC CTTTGCGACA | 540 |
| CATGGGGAAA CCTTCCCTCC CTTCTCTGT TGTACCAAT GGCCCTTAA GAGGAGCAGG | 600 |
| GCCACCTTGA AACTTGAGG ATATGGGGTA ACCCAGTGGG AGCGGGCAGG GAGGGCCCTT | 660 |
| GGAAACTGAC AGGGCTGGAG TATCTGCTG GGTTCAGCC CCGGTTCTG CAGGCACAGC | 720 |
| TGCCAGGCTC TCTGTTCAAC TTCCTGCCTC TGGTTTGCCC CGGCTCCCTC ACCCCCCTTA | 780 |
| CCCTGGAGTC CTTCTTCTA GGTGGGAGAT GCCATCCCAG CAGTGGAGGT GTTGAAGGG | 840 |
| GAGCCAGGGA ACAAGGTGAA CCTGGCAGAG CTGTTCAAGG GCAAGAAGGG TGTGCTGTTT | 900 |
| GGAGTTCCTG GGGCCTTCAC CCCTGGATGT TCCAAGGTGA GGCCCTTCCC CTTCTGAAGA | 960 |
| TCAGGACCTG GGGATCTTTT GTGTGCTCT TAAGTCCTCC ACATAGTCCT GATAGGACTC | 1020 |
| CTAAAAAGCA TTTCAGTGCC ATCACAAAAC AAGTAGAGCT GGGTAGAGCT GGGCGCGGTG | 1080 |
| GCTCACGCCT GTAATCCCAG CACTTTGGGA GGCCAAGGCG GGTGGATCAC GAGGTCAGGA | 1140 |
| GTCCAAAACC AGCCTGGCCA AGATGGTGAA ACCCTGTCTC TACTAAAAAT GCAAAAAAAT | 1200 |
| CAGCCGGATA TGGTGGCGGG CGCCTGTAAT CCCAGGTATT GGGGAGGCTG AGGCAGAGAA | 1260 |
| TTGCTTGAAC CCAGGAGGCG TAGGTTGCAG TGAGTGGAGA TCGTGCCTCT GCAGTCCAGC | 1320 |
| CTGGGTGAAA GAGCGAGACT CCGTCTCAA ATGAAAAAAA AAAAAGAAAA CAAGTAGAGA | 1380 |
| CTGCAAAAAG GGAACAGTAC CGGGAATGTT GGAGAAAAAC ATACTACAAT TAAATCCAAC | 1440 |
| ACCCCTGTTG GTCCTGCTAA ATGACAGGCA CTGTGGAAGG TGCTTGGGAC TCAGATAAAT | 1500 |
| AAGACAAAGA TCTGCCCATG GAAAGTTCAC GTCTGGACCA TAAGGCATTA GGTTCATTC | 1560 |
| TGAGCTTCCT AGTGGCCAAG GCAAAAAGGA AATAGAATGG TTTAGACAGC TCTCATTGTC | 1620 |
| TGATCAAAGG TGTGAGGCA GAGCACTGAG GAGGGCCTGG AGATAAAGGG TGGGCTGGGG | 1680 |
| GTCAGATGCA GTTATCCCTT TGCCGACCCT TTGTTCCCCT TCCTCAGACA CACCTGCCAG | 1740 |
| GGTTTGTGGA GCAGGCTGAG GCTCTGAAGG CCAAGGGAGT CCAGGTGGTG GCCTGTCTGA | 1800 |
| GTGTTAATGA TGCCTTTGTG ACTGGCGAGT GGGGCCGAGC CCACAAGGCG GAAGGCAAGG | 1860 |

| | |
|--|------|
| TGAGGTGAGG GGCCTGCAGG GAGTCAGGAC CAGGTAGGAT ATTCTTCTTG TGACCTCTAC | 1920 |
| TTTCTCTGCA GGTTCGGCTC CTGGCTGATC CCACTGGGGC CTTTGGGAAG GTGAGTGTTT | 1980 |
| CCCTGACCGC CACAGGGACA TGGCGGTGCG GGGAGCAGTG GGGGCCCTTG GCCTCTTCAA | 2040 |
| GGATTTCTGA CACTTTTCTC TGTCTCTTCT TAGGAGACAG ACTTATTACT AGATGATTCTG | 2100 |
| CTGGTGTCCA TCTTTGGGAA TCGACGTCTC AAGAGGTAAA AGTGGAGAGT CCTCTGTGGA | 2160 |
| GAAAGTCCTC TGTGGGAGAG AGTCCTCTGT GGGAGAGAGT CCTCTGTGGA GAGGGTCCTC | 2220 |
| TGTGGGAAGA GTCGTCTGTG GGGGAGATGT GTGGGAGAGA GTCCTGTGTG GGGAGAGTCT | 2280 |
| TCTGTAGGGG AGAGTCCTCT GGGGAGAGAG TCCTGTGTGG GGGAGAGTCC TCTGTGGGGA | 2340 |
| GAGTCCTCTG TGTGGAGAGA GTCCTGTGTG GTGGTGAGTC CTCTGTGGGG GAGAGTCCTC | 2400 |
| TGTGGGGGGA GTCCTCTCTG GAGTTCTCTT GGGCCCCTGG CTGTTCACTG CCTGTCTCCA | 2460 |
| TGCCCAGCCT CCAAGCCCAG GCTGATGCAG CTGGCTGGGC CCCTCTTTCC GGCAGGTTCT | 2520 |
| CCATGGTGGT ACAGGATGGC ATAGTGAAGG CCCTGAATGT GGAACCAGAT GGCACAGGCC | 2580 |
| TCACCTGCAG CCTGGCACCC AATATCATCT CACAGCTCTG AGGCCCTGGG CCAGATTACT | 2640 |
| TCCTCCACCC CTCCCTATCT CACCTGCCCA GCCCTGTGCT GGGGCCCTGC AATTGGAATG | 2700 |
| TTGGCCAGAT | 2710 |

(2) INFORMATION FOR SEQ ID NO: 11:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

GCCATCCCAG CAGTGGAGGT GTTTG

25

(2) INFORMATION FOR SEQ ID NO: 12:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

12

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

TTGAACAGCT CTGCCAGGTT CACC

24

(2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

TGGAGGTGTT TGAAGGGGAG CCAG

24

(2) INFORMATION FOR SEQ ID NO: 14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

CAGGTTACCC TTGTTCCCTG GCTC

24

(2) INFORMATION FOR SEQ ID NO: 15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

GGGTATGGGA CTAGCTGGCG

20

(2) INFORMATION FOR SEQ ID NO: 16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single

13

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

CTGGCCAACA TTCCAATTGC AG

22

(2) INFORMATION FOR SEQ ID NO: 17:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

ATGTTATGCA ACCCTTTGCG ACAC

24

(2) INFORMATION FOR SEQ ID NO: 18:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

GTGTTTGAAG GGGAGCCAGG GAAC

24

(2) INFORMATION FOR SEQ ID NO: 19:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

AGAGACAGGG TTTCACCATC TTGG

24

